

RESOLUTION AND SENSIBILITY INCREASE IN CR-39 PLASTIC COSMIC RAY DETECTOR

A. Laville, J. Pérez-Peraza¹, D. López², M. Balcazar-García³,
and A. López³

Instituto Nacional de Astrofísica, Óptica y Electrónica
Puebla, México

RESUMEN. Los rayos cósmicos son una parte más de la información que el Universo nos envía. De hecho es una valiosa información que no debemos desperdiciar. Una forma permanente de registrar esta información, son los SSNTD (detectores de estado sólido por trazas nucleares), en donde existe una búsqueda permanente para incrementar la resolución y sensibilidad de dichos detectores con el fin de perder la menor información posible. El detector plástico que proporciona actualmente la mejor resolución y sensibilidad es el polímero CR-39 (polímero del dietilenglicol bis alil carbonato). Un material similar (POLINAOE) ha sido desarrollado en el INAOE para el estudio de partículas solares y radiación cósmica galáctica. Se ha establecido de manera completamente empírica que la resolución y sensibilidad de estos materiales plásticos se modifica cuando el material es impurificado con ciertos aditivos, tales como el dioctil ftalato. En este trabajo se reportan resultados experimentales y su correspondiente interpretación en un intento de profundizar en el entendimiento de estos efectos, dentro del marco de incrementar la resolución y sensibilidad del material detector.

ABSTRACT. Cosmic rays are part of the information that the Universe send to us. As a matter of fact, it is a valuable information that we should not loose. A permanent way of registering this information is by using SSNTD (solid state nuclear track detectors) where a continuous search has been done for incrementing the resolution and sensibility of such detectors with the purpose of loosing the least possible information. At present the best resolution and sensibility has been obtained with a plastic detector known as CR-39 (polimer of dietylen glycol bis allyl carbonate). A similar detector material (POLINAOE) has been developed at the INAOE for studies of solar particles and galactic cosmis rays. It has been established in a completely empirical way that the resolution and sensibility of these detectors are modified when the material is doped with and additive such as dioctyl phtalate. Here we report experimental results and our interpretation, in an attempt to go deeper in the understanding of these effects, within the frame of our intentions to increase the resolution and sensibility of the detector material.

Key words: INSTRUMENTS — DETECTORS

1. On leave from Instituto de Geofísica, UNAM, 04510 México, D.F., México.
2. On leave from Universidad de Puebla, Facultad de Química, Puebla, Pue. México.
3. On leave from Instituto Nacional de Investigaciones Nucleares.

I. INTRODUCTION

In the world of astronomy a substantial quantity of information comes from cosmic rays. To detect cosmic rays, we need different kind of detectors than those used for light. A suitable kind of these detectors are the SSNTD, materials in which particles cause a damage while passing through them, damage that can be developed by etching technics, to see under the microscope the amplified particle track.

In nuclear track studies there is a constant search for improving the resolution and sensibility of the detector materials. For resolution we mean here the ability to distinguish between tracks of a given kind of particles of different but close energies, or particles of same energy but different mass, atomic number or effective charge. For sensibility we mean the capacity of the material to detect particles of enough low energy, that is that even particles of low energy produce a damage on the material because the ratio of etching track velocity to etching general or bulk velocity is high enough ($V_t/V_g \gg 1$). Since the particle energy loss (dE/dx) depends on parameters such as the mass A , the atomic number Z and the effective charge q^* , we can also refer resolution and sensibility for those other parameters.

It has been known for some time that as far as plastic detectors are concerned, when they are doped with some additives, the resolution and sensibility may change with respect to the plain (non doped) plastics. These are empirical results and the mechanism for explaining this change in resolution and sensibility has not yet been found. In the particular case of CR-39 it has been shown (Tarlé 1981) that when this is doped with dioctyl phthalate the resolution of the material increases with respect to that of CR-39 plain, though the sensibility slightly decreases; it has also been shown that the effects of phthalates and terephthalates are practically the same. In this work it is attempted to go deep in the understanding of the processes involved in the alterations of resolution and sensibility for nuclear track detection when a plastic is doped. At last instance it is searched how to increase both resolution and sensibility, to set limits for that improvement without changing drastically the physico-chemical properties of the material.

The present work is in some extent related with a previous work (Pérez-Peraza et al. 1983, 1984) where the main goal was to study the effect of finite temperature of the detector material on the nuclear track during particle energy deposition. Among the several conclusions of that work is that the control of both resolution and sensibility depend on both, the macroscopic and microscopic elasticity properties of the material. The former are represented through Young's modulus (Y) and the latter by the interatomic elastic constant (k_s), which in turn is directly proportional to the dissociation energy and inversely proportional to the square of the interatomic distance in the equilibrium position. Therefore, the natural suite is to search for experimental techniques allowing to control those macroscopic and microscopic parameters, and in turn to get control on resolution and sensibility. We know from several works that dopants are able to modify both resolution and sensibility so, in some way they must act on the macroscopic and microscopic parameters, as we will hypothesize in the next section.

In the search for understanding the intercorrelations between the doping techniques and the macroscopic and microscopic parameters, we proposed in a preliminar work (Laville et al. 1985) that additives like phthalates and terephthalates act either as polymer chain openers or as blockers of cross-linking, whereas additives of the kind of benzoates may act only as blockers (fig. 1).

In that work, it was established that benzoate additives do not increase the resolution as high as phthalates or terephthalates, but it was better than the resolution without additives in CR-39 plain. Also it was found that the sensibility did not decrease with benzoate as much as doped with phthalates or terephthalates. We concluded that the predominant effect in increasing resolution was chain opening.




Fig. 1. Reaction sites in phthalates, terephthalates and benzoates.

II. THEORETICAL APPROACH

Our depart thesis in relation with the conclusions on the previous work, is that the Young modulus may be increased by increasing the separation between polymeric chains, that is, making softer the reticulation; or may be decreased by closing the separation between chains, that is, making the reticulation more compact. On the other hand, the interatomic spring constant may be increased or decreased by increasing or decreasing the cohesion energy respectively. The increase or decrease of the macroscopic elasticity does not necessarily entail the increase or decrease of the microscopic elasticity and conversely. How do these processes are translated on resolution and sensibility of the material, to nuclear tracks of penetrating projectiles? If separation between chains is increased, the interaction cross section between projectile and target atoms diminishes, the energy loss of projectiles is lowered and the particle range increases leaving a longer track. This entails an increase in resolution with respect to a more compact reticulation, because it is easier to discriminate between tracks of particles of neighbor energies as the reticulation becomes softer and bond density decreases: let us consider two particles of energies E_i , E_j above the threshold energy, such that $E_i - E_j = \Delta E$. Once particle of energy E_j has been stopped, it is easier for particle of energy E_i to cause still further damage on the plastic before stopping, on a softer net, whereas in a more compact reticulation the bond density is higher and so, for the same energy difference ΔE , particle energy E_i is stopped very close to that of energy E_j , causing similar damage. In other terms with the separation of chains, damage along the track increases and this causes that tracks become more distinguishable among them. Under this point of view, resolution decreases by compacting the reticulation (fig. 2).

Now concerning sensibility, if separation between polymeric chains is increased, there is a compromise between, for one side, the increase in sensibility because of the decrease of dissociation energy of atomic bonds due to the enlargement of the reticulation cells with the subsequent decrease of cohesion (the chain net becomes softer, such that particles of lower energy are above the threshold energy) and from the other side, the decrease in sensibility by the fact that the cross section decreases with

cell enlargement. However this compromise is solved in the case of additives that contain a benzoic ring (as it is the case of the previously mentioned) because the benzoic ring chemical structure has a high stability that creates a bigger energy of cohesion, that increases substantially the dissociation energy in the sites of reaction between additives and monomeric molecules. The result is a decrease in sensibility as established by Tarlé (1981). Therefore projectiles of energy lower than a certain threshold value are not able anymore to cause a damage in the doped plastic as they do in the plain one. However once they have the threshold energy they are better resolved in a doped plastic.





Fig 2.- Effect of the softening in the reticulation.

Now according to our thesis, four different things can happen during polymerization when we add the additives. Accordingly they are:


1.- OPENING: When molecules of the additives have two reaction sites as it can be the case of phthalates and terephthalates (see fig. 1) they interpose between the monomer molecules of two different chains, reacting with them on both sites during crosslinking of the monomer molecules thus having as a result the opening between chains and the softening of the cells, otherwise would have become more compact by the single effect of crosslinking without the additive (fig. 3).

2.- BLOCKING: Blocking of the chains is expected when the additive molecule have only one reaction site (see fig. 1), for instance benzoates, in which case the molecule reacting with one free radical of a monomer chain will avoid that this particular chain will undergo crosslinking with another chain in that particular reaction site. The result is that the cohesion of the net becomes softer in those blocked sites. This is translated in a density decrease of the reticulation in relation with the non doped plastic, this is, though the cells become larger they are not so open or soft as in the case of the two reaction sites additives (fig. 4). We have to remember that phthalates and terephthalates can also work as blockers, when they present only one reaction site.



M=Monomer , D=Dopant.

Fig 3.- Opening hypothesis.




M=Monomer , D=Dopant.

Fig 4.- Blocking hypothesis.

3.-RETICULATION: Suppose now that during polimerization instead of a dopant molecule, an individual monomeric molecule react with two monomeric chains causing crosslinking between them. The result is that the reticulation cells in those particular crosslinking sites becomes larger even when crosslinking occurs by means of a dopant molecule, because of the monomer molecules being physically bigger than the dopant molecules (fig. 5). Remember that this can happen even if no dopant is added.

4.- FILLING: Our last hypothesis supposes that now the dopant molecules place themselves between monomeric chains but without reacting with them. Since in this case the dopant molecule is not broken, the dopant molecule is larger than the monomeric molecules Therefore during conformation of the reticulation, the cells in those places becomes larger (fig. 6). We have to notice that benzoates can be smaller than the monomeric molecules as it is our particular case with benzyl benzoate, unexplaining how by this hypothesis, resolution can be increased with this particular benzoate.




Plain.

Doped.

M=Monomer , D=Dopant.

Fig 5.- Reticulation theory.



Plain.

Doped.

M=Monomer , D=Dopant

Fig 6.- Filling theory.

III. EXPERIMENTAL PROCEDURE

1.-ADDITIVES AND POLYMERS: Three different polymer sheets 5x5 cm.² and 380 μ m. thick were made. One plain and two doped. The two dopants used were dioctyl phtalate (DOP) and benzyl benzoate (BB). All three sheets were polymerized using benzoyl peroxide (BP) as initiator in an oven at a constant temperature of 70 °C and a curing period of 72 hours (Table I.)

TABLE I.- Polymer sheets 5x5 cm.² 380 μ m. thick.

Polymer name	Initiator	Initiator conc.	Dopant	Dopant conc.
POLINAOE	BP	3%	-	-
POLINAOE(DOP)	BP	3%	DOP	3%
POLINAOE(BB)	BP	3%	BB	3%

NOTE: Initiator and dopant concentration is percentage by weight.

2.-IRRADIATION: Of the 5 cm.² sheets, seven squares of 1 cm.² of each plastic were cutted and irradiated with particles from a ²⁵²Cf and ²⁴¹Am radioactive sources. The samples were placed in an irradiation fan. The fan had several slots perpendicular to the irradiation channels so that absorbers may be interposed in the particles trajectories before they could hit the plastics, and so different energies could be obtained. The absorbers where previously calibrated to give us the energy of the particle hitting the detector. The 21 samples were identified according to their additive and the irradiation they were going to receive. Once the samples and the absorbers were placed in the irradiation fan it was assembled in the versatile irradiation chamber (Balcazar-García and Pineda 1983) where they were irradiated with α particles.

3.-ETCHING AND READING: After the samples were irradiated these were etched in a NaOH solution 6.25 N at 70°C until the tracks diameters were measurable. This happened at an etching time of 3.5 hours. Fifty diameters were measured for each sample and the statistical media was calculated. The same procedure was repeated for 5.5, 7.5, 9.5 and 11.5 hours, etching times. In addition one of the seven samples of each kind of plastic was irradiated with fission fragment from the Cf source. One fission fragment track was selected in each plastic and identified perfectly by a nearby marking. In this way we can track the evolution of the fission track diameter in time for the several etching times. It is well known that the slope of a graph of track diameter of the fission fragment against etching time is the bulk etching velocity (V_g).

IV. DATA ANALYSIS

Track diameters as a function of energy behave exponentially (Balcazar-García and Cuauhtecat1 1983):

$$\phi = k_1 \exp(-k_2 E)$$

k_1 and k_2 are characteristic of each etching time and for each type of plastic. To evaluate them is necessary to graph the data for just one etching time, in a semilogarithmic scale. Here we evaluate the constants for etching times 7.5 and 9.5 hours for the whole set of samples. Now, knowing the empirical expression between track diameter ϕ and energy E , we want to relate with V_t . Using $V = V_t / V_g$ and the well known expression:

$$\phi = 2 V_g t \sqrt{(v+1)/(v-1)}.$$

we obtain:

$$V = k_1^2 \exp(-2k_2 E) + 2V_t / 2V_t - k_1^2 \exp(-2k_2 E).$$

where t is the correspondent etching time for which k_1 and k_2 were calculated.

from the latter expression we can plot V against E and from the fact that $V = V_t / V_g$ we can plot V_t against E for the 7.5 and 9.5 etching times. From figures 7,8,9,10, we clearly see the difference in slopes for the three plastics. We have to note here that as steeper is the slope of these graphs, the better the resolution of the plastic in question i.e. for small changes in energy bigger changes in V or V_t for steeper slopes. Then it is seen that the one with more resolution is the doped with DOP, then the doped with BB and at last the plain one.

V. DISCUSSION AND CONCLUSIONS

As we mentioned earlier, the increment in resolution may happen by one of the four hypothesis or by a combination of them. From figures 7,8,9,10, we discard the reticulation hypothesis because if true, we would expected that the plain plastic would have more resolution than the other ones, as this kind of reticulation happens in the absence of additives. Now, if we notice that plastic doped with DOP increases the resolution better than the one doped with BB and the latter one increases has a better resolution than the plain plastic, then if the filling theory were true, the BB additive should not have effect at all because of its molecules being smaller than the monomer molecules, filling nothing at all. The preceeding reasoning discards the filling theory. The difference in resolution between DOP and BB, confirms our earlier result that DOP works mainly as an opener but that it can work in a lesser extent, also as a blocker, this is shown by the improved resolution of the plastic doped with BB compared to the plain one

and considering that this additive can only work as a blocker after discarding the reticulation and filling theories.

From the careful analysis of the four hypothesis and the results shown, we can conclude that the most effective dopants are the ones that act mainly as openers as DOP does. We suggest the search for dopants that can open the chains more to corroborate completely this theory and at the same time, this new dopants should be absent of the benzoic ring, so that sensibility is not decreased.

As a last word, reticulation and filling theories are highly improbable especially the reticulation theory as it opposes completely to the other three hypothesis and does not check at all with the results presented here and on our earlier work.




Fig 7.- V against E (7.5 hours).




Fig 8.- V against E (9.5 hours).




Fig 9.- V against E (7.5 hours).




Fig 10.- V against E (9.5 hours).

REFERENCES

- Balcazar-García, M. and Cuauhtecatl, V.; "The Nucleus" ; 20; 129;1983.
 Balcazar-García, M. and Pineda, H.; "The Nucleus" ; 20;75;1983.
 Laville A., Pérez-Peraza J., Alvarez M., and Estrada M.R.; Proc. of 19th Int. Cosmic Ray Conf.; 2; 29; 1985.
 Pérez-Peraza J., Laville A., and Balcazar-Garcia M.; Proc. of 18th Int. Cosmic Ray Conf.; 18; 11; 1983.
 Pérez-Peraza J., Laville A., Balcazar-Garcia M., and Camacho S.; Nuclear tracks and rad. measurements; 8; 147; 1984.
 Tarlé G.; Proc. of 17th Int. Cosmic Ray Conf.; T2-13; 1981.

M. Balcazar-García, A. Laville, A. López, D. López, and J. Pérez-Peraza: Instituto Nacional de Astrofísica, Óptica y Electrónica, Tonantzintla, Apartados Postales 51 y 216, 72000 Puebla, México.

INDICE DE AUTORES/AUTHORS' INDEX

- Abad, C., *Véase* Stock y Abad.
- Abraham, Z. e A.C.O. Cancoro, Modêlos teóricos de linhas de recombinação em radio frequências para regiões H II, 603.
- Abraham, Z., E. Scalise Jr., L.C.L. Botti, A.C.O. Cancoro, J.L.M. do Vale, L.F. del Ciampo, C.E. Tateyama, J.W.S. Vilas Boas, J.L. Homor, e P. Kaufmann, Observações do Cometa de Halley no contínuo de 22 GHz e 44 GHz, e na raia maser de vapor de H₂O em 22.2 GHz, 671.
- Abraham, Z. *Vea-se* Botti e Abraham.
- Acosta J.A., M. Kidger, M. Prieto, J. Cepa, and C. Muñoz Tuñón Maps of the Emission of CN and other Molecular Species in the Spectrum of Comet Halley, 644.
- Aguilar, L.A., J.P. Ostriker, and P. Hut, Destruction Mechanisms for the Globular Clusters in the Milky Way, 277.
- Alcalá, J.M. and A. Arellano Ferro, The Distance and Reddening of the Open Cluster NGC 7790 and the Luminosity of its Three Cepheids, 440.
- Allen, C., M.A. Martos, and A. Poveda, The Galactic Orbit of the Remarkable High-Velocity Wide Binary LDS 519, 213.
- Allen, C. and M.A. Martos, The Galactic Orbits of Nearby, High-Velocity Stars, 224.
- Allen, C. and M.A. Martos, A Simple Realistic Model of the Galactic Mass Distribution for Orbit Computations, 226.
- Alvarez, M., D. Ballereau, J.P. Sareyan, J. Chauville, R. Michel, and J.M. Le Contel, Spectroscopy and Photometry of some Intrinsic B Variable Stars, 315.
- Alvarez-Madrigal M., *See* Pérez-Peraza *et al.*
- Alvarez-Madrigal M., *See* Pérez-Peraza *et al.*
- Andrade, A.A., S.M. Viegas-Aldrovandi, R.B. Gruenwald, Balmer Lines in Active Galactic Nuclei, 90.
- Andrade, A.A., S.M. Viegas-Aldrovandi, and R.B. Gruenwald, Balmer Lines in Active Galactic Nuclei, 90.
- Angeles, F., Sistema de adquisición de datos para un fotómetro doble basado en una microcomputadora, 749.
- Angeles, F., E. Carrasco, S. Cuevas, R. Enríquez, L. Gutiérrez, R. Langerica, E. Ruiz y E. Sacristán, Guiador excéntrico para el telescopio de 2.1 mts. en San Pedro Mártir, B.C., 736.
- Antonelli, P., *Véase* Chapellier *et al.*
- Arellano Ferro, A., L. Parrao, A. Giménez, M. López Arroyo, and D.F. Sasselov, H α Profile Variations in Epsilon Aurigae, 418.
- Arellano Ferro, A., *See* Alcalá and Arellano Ferro.
- Arellano Ferro, A., *See* Evans and Arellano Ferro.
- Arnal, M., H. Levato, B. García, and N. Morrell, The Open Cluster NGC 6193: Spectral Types, Axial Rotation and Binarity, 423.
- Arnal, E.M., *See* Cappa de Nicolau *et al.*
- Arnal, E.M., *See* Silva *et al.*
- Auer, L.H., *See* Koenigsberger and Auer.
- Auer, L.H., *See* Koenigsberger and Auer.
- Bajaja, E., C. Cappa de Nicolau, M.C. Martín, R. Morras, C.A. Olano, y W.G.L. Pöppel, Estudio de algunas nubes de alta velocidad del hemisferio sur, 604.
- Bajaja, E., *Véase* Silva *et al.*
- Balcazar-García, M., *See* Laville *et al.*
- Ballereau, D., *See* Alvarez *et al.*
- Balthazar, J.M., J.L. Sagnier, S.F. Mello, J. Koiller, e T. Yokoyama, Sobre algumas técnicas de perturbação, utilizadas no problema ressonante 3/1, 639.
- Balthazar, J.M., *See* Yokoyama *et al.*
- Barral, J.F., G.F. Bisiacchi, C. Firmani, G. Koenigsberger, and J. Wampler, Spectrophotometry of Northern Wolf-Rayet Stars: A Sample of WN4, WN5, and WN6 Stars, 284.
- Barral, J.F., M. Ríos Berumen, M. Ríos Herrera, A. Quintero, A. Pani, R. Peniche, and J.H. Peña, Earth Based Observations of Comet Halley, 660.
- Barral, J.F., *See* Serrano P.G. *et al.*
- Barrera, L.H., and N. Vogt, Spectroscopic Monitoring of Southern Be Stars, 323.
- Beck, R., *See* Loiseau *et al.*
- Beckman, J., J. Cepa, M. Prieto, and C. Muñoz Tuñón, NGC 2403: A Flocculent Galaxy with Two Principal Centres of Star Formation, 134.
- Beckman, J., *See* Muñoz Tuñón *et al.*
- Benevides-Soares, P., Representation of Systematic Differences Astrolabe-Catalog, 454.
- Benevides-Soares, P., *See* Rossi and Benevides-Soares.
- Bergmann, T.S. and M.G. Pastoriza, Galaxies with Strong Nitrogen Lines, 106.
- Bergmann, T.S., *See* Ducati *et al.*
- Bevilacqua, C.M., *See* Ducati *et al.*
- Bevilacqua, J.S. and W. Sessin, First Order Perturbation in the Enceladus-Dione System, 627.
- Bica, E., *See* Dottori *et al.*
- Bisiacchi, G.F. y C. Firmani, Deposición de un fotocátodo para uso astronómico, 771.
- Bisiacchi, G.F., *See* Barral *et al.*
- Bisiacchi, G.F., *See* Firmani and Bisiacchi.
- Bodenheimer, P., *See* Franco *et al.*
- Bodenheimer, P., *See* Tenorio-Tagle *et al.*
- Bohigas, J., and J. González, The Galactic Thermal Pressure Gradient, 228.
- Bonatto, C., *See* Ducati *et al.*
- Boriakoff, V., *See* Silva *et al.*
- Botti, A.F.L., and J.R.D. Lépine, Study of the Galactic Structure using Different Objects as "Tracers", 225.
- Botti, L.C.L., and Z. Abraham, Low Term Variability of Radio Sources in the Frequencies of 22 GHz and 44 GHz, 97.
- Botti, L.C.L., *Vea-se* Abraham, Z. *et al.*
- Brandi, E., O.E. Ferrer, and J. Sahade, Line Structures in the IUE Spectra of γ_2 Velorum, 300.
- Bravo, S., B. Mendoza, R. Pérez-Enríquez y J. Valdés-Galicia, Actividad de los hoyos coronales solares de baja latitud, 688.
- Bruzual, G., Is there any Evidence of Evolution in the Color Distribution of Galaxies from J = 13 to J = 24 Magnitude?, 169.
- Bruzual, A., *See* Magris and Bruzual.
- Bulka, P., *Véase* Ferrín *et al.*
- Bulka, P., *Véase* Naranjo *et al.*
- Burbidge, G., Continuity in the Violent Phenomena Taking Place in the Nuclei of Galaxies and QSOs, 81.
- Calvet, N., and Z. Marín, The Photosphere of T Tauri Stars, 353.
- Canal, R., J. Isern, and J. Labay, Type I Supernova Models, 245.
- Canal, R., *See* Isern, *et al.*
- Cancoro, A.C.O., *Vea-se* Abraham e Cancoro.
- Cancoro, A.C.O., *Vea-se* Abraham *et al.*
- Cantó, J., *See* Curiel *et al.*

- Cappa de Nicolau, C., V.S. Niemela, G. Dubner, and E.M. Arnal, RCW114: Another Case of H I Bubble Related to a WR Star, 611.
- Cappa de Nicolau, C., *Véase Bajaja et al.*
- Carballo, R., A. Mampaso, and N. Epchtein, Search for the Near Infrared Counterparts of Compact H II Regions and/or H₂O Masers, 485.
- Cardona, O., E. Chavira, I. Furenlid, y B. Iriarte, El sistema CCD de Tonantzintla. Pruebas y planes futuros, 744.
- Cardona, O., *See Iriarte et al.*
- Carpenter, K.G., *Véase Hagen et al.*
- Carral, P., W.J. Welch, and M.C.H. Wright, HCO⁺ and SO Emission Associated with the Region G34.3+0.2, 506.
- Carrasco, E., *Véase Angeles et al.*
- Carrasco, L., R. Costero, and R. Stalio, Resonance Line Profiles in Galactic Disk UV-Bright Stars, 301.
- Carrasco, L., I. Cruz-González, and G.L. Grasdalen, On the Masses of the Circumstellar Dust Disks around the YSO's R Mon and R50, 518.
- Carrasco, L., A. Chelli, H. Zihnecker, I. Cruz-González, and C. Perrier, Fast Scanning Slit Near IR Photometry of T Tauri Stars in Close Binaries: Elias 22 and Chamaleon I, 359.
- Carrasco, L., *See Chelli et al.*
- Carrasco, L., *Véase Cruz-González et al.*
- Carrasco, L., *See Cruz-González et al.*
- Carrasco, L., *Véase Navarro et al.*
- Carrasco, L., *See Recillas-Cruz et al.*
- Carrasco, L., *See Tapia et al.*
- Cavalcanti, R.L., *See Ducati et al.*
- Cepa, J., *See Acosta et al.*
- Cepa, J., *See Beckman et al.*
- Cersosimo, J.C., *Véase Silva et al.*
- Cervantes, A.M., *See Pişmiş and Cervantes.*
- Cieslinski, D., *See Steiner et al.*
- Clariá, J.J., R.F. Sisteró, y E. Lapasset, Observaciones fotométricas en bandas moleculares del Cometa Halley, 651.
- Clariá, J.J., *See Lapasset et al.*
- Clauzet, L.B.F., S. Débarbat, and F. Chollet, Optical and Radio Positions of α Scorpii, 455.
- Clauzet, L.B.F., and M.E. Mattos, Provisional Values of Internal Group Corrections for the VL3 Astrolabe Catalogue, 453.
- Clocchiatti, A. and H.G. Marraco, The Small Scale Interstellar Dust Distribution, 613.
- Cobos, F.J., y M.J. Galán, Análisis de alteraciones en la imagen debidas a descolimación de un telescopio, 768.
- Cobos, F.J., y M.J. Galán, Sistemas correctores de campo para el telescopio Ritchey-Chretien UNAM212, 770.
- Cobos, F.J., y R. Langarica, Integración del guiador excéntrico al telescopio UNAM212 y su repercusión en las imágenes obtenidas con diversos instrumentos, 739.
- Cobos, F.J., y C. Tejada de Vargas, Calidad de imagen del telescopio UNAM212, 766.
- Cobos, F.J., *Véase Galán y Cobos.*
- Cobos, F.J., *Véase Galán y Cobos.*
- Colomb, F.R., *Véase Silva et al.*
- Contini, M., *See Viegas-Aldovandi and Contini.*
- Copetti, M.V., *See Dottori and Copetti.*
- Copetti, M.V.F., y H.A. Dottori, Fotometría W_H β y [OIII]/H β de regiones H II y la historia de la formación estelar reciente en la Pequeña Nube de Magallanes, 191.
- Corona, M.G., A Hydrodynamical Three-Component Model of the Universe during the Recombination Era, 52.
- Corral, L. and G. Koenigsberger, Photometric Observations of the Be/X-Ray Binary System 2S0114+650 = LSI+65°010, 330.
- Cortés, G., *See Gutiérrez-Moreno et al.*
- Cortés, P., and F. Sala, A Model of the Stellar Velocity Field, 206.
- Costa, E., *See MacConnell et al.*
- Costa E., *See Wing et al.*
- Costa, R.D.D., *See Ducati et al.*
- Costero, R., *See Carrasco et al.*
- Costero, R., *Véase Navarro et al.*
- Cruz-González, I., L. Carrasco, y M.P. Ramos, Fuentes infrarrojas a 2.2 μ m en la región asociada al glóbulo Ori-I-2, 519.
- Cruz-González, I., Recillas-Cruz, E., and Carrasco, L., Near Infrared Maps of Elliptical and S0 Galaxies Detected by IRAS, 157.
- Cruz-González, I., *See Carrasco et al.*
- Cruz-González, I., *See Carrasco et al.*
- Cruz-González, I., *See Chelli et al.*
- Cruz-González, I., *See Recillas-Cruz et al.*
- Cuevas, S., Un interferómetro de rotación de frente de onda con modulación, 753.
- Cuevas, S., *Véase Angeles et al.*
- Cuevas, S., *Véase Tinoco y Cuevas.*
- Curiel, S., J. Cantó, and L.F. Rodríguez, A Model for the Thermal Radio Continuum Emission Produced by a Shock Wave and its Application to the Herbig-Haro Objects 1 and 2, 595.
- Chapellier, E., J.M. Le Contel, J.C. Valtier, S. González-Bedolla, D. Ducatel, P.J. Morel, J.P. Sareyan, I. Geiger y P. Antonelli, Variaciones fotométricas en la estrella B Iota Herculis, 424.
- Chapellier, E., *Véase González et al.*
- Chatterjee, T.K., A Dynamical Explanation for the Low Frequency of Ring Galaxies, 120.
- Chauville, J., *See Alvarez et al.*
- Chavarría-K., C., J. Ocegueda, E. de Lara, U. Finkenzeller, and E.E. Mendoza, An Observational Study of the Herbig Ae Star VV Serpentis and R-Stars Associated with its Dark Cloud, 338.
- Chavira, E., The P/Halley's Comet Encountering the Interplanetary Medium, 678.
- Chavira E., *Véase Cardona et al.*
- Chavira, E., *See Iriarte et al.*
- Chelli, A., Comparison Between Image Plane Phase Reconstruction Methods in Optical Interferometry, 751.
- Chelli, A., C. Perrier, I. Cruz-González, and L. Carrasco, High Spatial Resolution IR Observations and Variability of the Nuclear Region of NGC 1068: Structure and Nature of the Inner 100 Parsec, 102.
- Chelli, A., *See Carrasco et al.*
- Chollet, F., *See Clauzet et al.*
- Danks, A.C, Carbon Chemistry of the Interstellar Medium, 614.
- Débarbat, S., *See Clauzet et al.*
- del Ciampo L.F., *Véase Abraham et al.*
- de Lara, E., *See Chavarría-K. et al.*
- de Lara, E., *See López and de Lara.*
- de la Reza, R., *See Quast et al.*
- de la Reza, R., *See Torres et al.*
- Delgado, A., *Véase González et al.*
- Dottori, H.A., Evolution of H II Regions, 463.
- Dottori, H.A. and M.V. Copetti, Spiral Structure and Age Distribution of the H II Regions in the Milky Way, 223.
- Dottori, H.A., J. Melnick, and E. Bica, NGC 2209: The Nature of the Dark Patch through the HR Diagram, 183.
- Dottori, H.A., *Véase Copetti and Dottori.*
- Dottori, H.A., *See Ducati et al.*
- Dottori, H.A., *See Schmidt and Dottori.*
- Dottori, H.A., *See Schmidt et al.*
- Dottori, H.A., *See Zepka and Dottori.*
- do Vale, J.L.M., *Véase Abraham et al.*
- Dubner, G., *See Cappa de Nicolau et al.*
- Ducatel, D., *Véase Chapellier et al.*
- Ducati, J.R., T.S. Bergmann, C.M. Bevilacqua, C. Bonatto, R.L. Cavalcanti, R.D.D. Costa, H.A. Dottori, L. Girardi, D. Hadjimech, S.O. Kepler, S.H.B. Livi, M.G. Pastoriza, J.F. Santos, A. Schmidt and M.F.S. Schroder, Strip Photometry of Comet Halley, 664.
- Dultzin-Hacyan, D. Low Ionization Emission Lines in Quasars, 94.

- Enríquez, R., *Véase* Angeles *et al.*
 Epchtein, N., T. Lébertre, J.R.D. Lépine, P. Marques dos Santos, O.T. Matsuura, and E. Picazzio, The Energy Distribution of 630 Sources from the Valinhos 2 μ m Survey, 375.
 Epchtein, N., *See* Carballo *et al.*
 Evans, N.R., and A. Arellano Ferro, The Luminosities of the Binary Cepheids SU Cyg, SU Cas and W Sgr, 435.
 Falco, E.E., Ultimas noticias sobre el sistema de lente gravitacional 0957+561, 127.
 Feinstein, C., *Véase* Forte *et al.*
 Ferrer, O.E., *See* Brandi *et al.*
 Ferrín, I., F. Fuenmayor, O. Naranjo, P. Bulka, y C. Mendoza, Procesamiento digital de imágenes del Cometa Halley, 669.
 Ferrín, I., *See* Naranjo *et al.*
 Fierro, J., *See* Peimbert *et al.*
 Finkenzeller, U., *See* Chavarría-K. *et al.*
 Firmani, C., and G.F. Bisiacchi, Star Formation in Blue Compact Galaxies, 171.
 Firmani, C., *See* Barral *et al.*
 Firmani, C., *Véase* Bisiacchi y Firmani.
 Flores Gutiérrez, J.D., Eventos planetarios periodicos, 643.
 Forte, J.C., E.I. Vega, M. Méndez, y C. Feinstein, Morfología CCD de Galaxias Activas: NGC 7552, 104.
 Franco, J., Asamblea general de la V Reunión Regional Latinoamericana de la Unión Astronómica Internacional, 16.
 Franco, J., G. Tenorio-Tagle, P. Bodenheimer, M. Rozyczka, and I.F. Mirabel, The Origin of the Orion and Monoceros Molecular Cloud Complexes, 240.
 Franco, J., *See* Tenorio-Tagle *et al.*
 Fricke, K.J., *See* Klapp, *et al.*
 Fuenmayor, F.J., The C/M5⁺ Ratio in the Galactic Disc, 379.
 Fuenmayor, F., *Véase* Ferrín *et al.*
 Fuenmayor, F., *Véase* Naranjo *et al.*
 Furenlid, I., *Véase* Cardona *et al.*
 Galán, M.J., y F.J. Cobos, Sistemas correctores de campo para el telescopio cassegrain IAC80, 767.
 Galán, M.J., y F.J. Cobos, Determinación de características ópticas del telescopio OAN150, 769.
 Galán, M.J., *Véase* Cobos y Galán.
 Galán, M.J., *Véase* Cobos y Galán.
 Galindo Trejo, J., Collisionless Tearing Instability as the Cause of an observed Structure in the Plasma Tail of Comet Halley, 672.
 Gallegos, A. and J. Pérez-Peraza, Determination of Energy Spectra of Solar Electrons under Different Scenarios in Solar Flare Sources, 700.
 Gallegos-Cruz, A., *See* Pérez-Peraza and Gallegos-Cruz.
 Garay, G., The Orion Radio Zoo: Pigs, Deers and Foxes, 489.
 García, B., *See* Arnal *et al.*
 García, B., *See* Levato *et al.*
 García, B., *See* Levato *et al.*
 García-Barreto, J.A., *See* Gómez *et al.*
 García-Berro, E., *See* Isern *et al.*
 Garrido, R., T. Gómez, R. Peniche, and J.H. Peña, LZ Her: An Eclipsing Binary Star, 416.
 Garrido, R., *See* González *et al.*
 Garrido, R., *Véase* González *et al.*
 Geiger, I., *Véase* Chapellier *et al.*
 Giménez, A., *See* Arellano Ferro *et al.*
 Giménez, A., *See* González *et al.*
 Girardi, L., *See* Ducati *et al.*
 Giridhar, S., Spectroscopic Studies of the Southern Cepheid β Dor, 431.
 Gómez, A.E., and J. Torra, Hipparcos. The Input Catalogue, 441.
 Gómez, D.O., and V.S. Niemela, A Spectrographic Study of Two Of Stars HD163758 and HD117797, 293.
 Gómez, M.N., *See* Lapasset *et al.*
 Gómez, T., *See* Garrido *et al.*
 Gómez, Y., L.F. Rodríguez, and J.A. García-Barreto, VLA Observations of Twice Ionized Helium in NGC 6302, 560.
 González, A., *See* Serrano *et al.*
 González, J., *See* Bohigas and González.
 González, S., J.P. Sareyan, R. Garrido, A. Delgado, y E. Chapellier, Variación en la amplitud de la curva de luz de la variable del tipo β CMA, δ Cet, 391.
 González, S.F., A. Rolland, A. Giménez, P. López de Coca, R. Garrido, M.A. Hobart, and J.H. Peña, Photometric Variability of the Binary HD 1826, 410.
 González, S.F., *See* Peña *et al.*
 González-Bedolla, S., *Véase* Chapellier *et al.*
 Grasdalen, G.L., *See* Carrasco *et al.*
 Gruenwald, R.B., *See* Viegas-Aldovandi and Gruenwald.
 Gruenwald, R.B., *Véase* Andrade *et al.*
 Gruenwald, R.B., *See* Andrade *et al.*
 Gutiérrez, L., *Véase* Angeles *et al.*
 Gutiérrez-Moreno, A., H. Moreno, and G. Cortés, PC 11: Sym-biotic Star or Planetary Nebula?, 344.
 Gutiérrez-Moreno, A., *See* Moreno *et al.*
 Hacyan, S., *See* Herrera and Hacyan.
 Hadjimichef, D., *See* Ducati *et al.*
 Hagen, W., K.G. Carpenter, y R.E. Stencel, Observaciones con IUE e IRAS de gigantes y supergigantes rojas, 380.
 Hasegawa, T., *See* Peimbert *et al.*
 Hasse I., *See* Pısmiř *et al.*
 Hernández, C., *See* Levato *et al.*
 Hernanz, M., *See* Isern *et al.*
 Herrera, M.A., and S. Hacyan, Thermal Relaxation Time of a Mixture of Relativistic Electrons and Neutrinos, 47.
 Hobart, M.A., *See* González *et al.*
 Hobart, M.A., *See* Peña *et al.*
 Hobart, M.A., *See* Rolland *et al.*
 Homor, J.L., *Véase* Abraham *et al.*
 Hummel, E., *See* Loiseau *et al.*
 Hut, P., *See* Aguilar *et al.*
 Ibañez S., M.H., Radiation Hydrodynamics in Astrophysics, 573.
 Iriarte, B., E. Chavira, and O. Cardona, Changes in the Be Stars in a Cassiopeia-Perseus Region, 336.
 Iriarte, B., *Véase* Cardona *et al.*
 Isern, J., R. Canal, E. García-Berro, M. Hernanz, and J. Labay, Collapse of White Dwarfs in Low Mass Binary Systems, 260.
 Isern, J., R. López, and E. Simonneau, Radiative Transfer in Type I Supernovae Atmospheres, 252.
 Isern, J., *See* Canal *et al.*
 Isobe, S., Some Developments in Optical Telescope Instrumentation and Data Reduction Systems in Japan, 722.
 Jablonski, F.J., *See* Steiner *et al.*
 Jugaku, J., *See* Peimbert *et al.*
 Kang, H., *See* Shapiro and Kang.
 Kaufmann, P., *Véase* Abraham *et al.*
 Kepler, S.O., *See* Ducati *et al.*
 Kidger, M., *See* Acosta *et al.*
 Kirhakos, D.S., *Véase* Steiner e Kirhakos.
 Klapp, J., N. Langer, and K.J. Fricke, Uncertainties in the Determination for the Upper Mass Limit for Zero-Age Main Sequence Stars, 265.
 Koenigsberger, G., and L.H. Auer, UV Line Profile Variations in Wolf-Rayet Stars, 271.
 Koenigsberger, G., and L.H. Auer, Identifying Lines in the IUE Spectrum of the Wolf-Rayet Star HD193077, 277.
 Koenigsberger, G., *See* Barral *et al.*
 Koenigsberger, G., *See* Corral and Koenigsberger.
 Koiler, J., *Véase* Balthazar *et al.*
 Koiler, J., *Véase* Balthazar *et al.*
 Kondo, Y., *See* Solivella *et al.*

- Labay, J., *See Canal et al.*
 Labay, J., *See Isern et al.*
 Lagua, R.O., and S.M. Viegas-Aldrovandi, Dust Heating by Ultra-violet Accretion Disk Radiation, 107.
 Lahulla, J.F., Fotometría UBVR_I de las regiones S61, S129, S285 y S299, 437.
 Langarica, R., *Véase Angeles et al.*
 Langarica, R., *Véase Cobos y Langarica.*
 Langer, N., *See Klapp, et al.*
 Lappaset, E., *Véase Clariá et al.*
 Lapasset, E., M.N. Gómez, and J.J. Clariá, A Photometric Analysis of the Massive Contact Binary BR Muscae, 402.
 Laville, A., J. Pérez-Peraza, D. López, M. Balcazar-García, and A. López, Resolution and Sensibility Increase in CR-39 Plastic Cosmic Ray Detector, 776.
 Lazzaro, D., *See Vieira Martins et al.*
 LéBertre, T., *See Epchtein et al.*
 LeContel, J.M., *See Alvarez et al.*
 LeContel, J.M., *Véase Chapellier et al.*
 Lépine, J.R.D., *See Botti and Lépine.*
 Lépine, J.R.D., *See Epchtein et al.*
 Levato, H., C. Hernández, N. Morrell, and B. García, Binary Frequencies in the Open Cluster IC 2602, 414.
 Levato, H., S. Malaroda, B. García, and N. Morrell, Spectroscopic Binary Frequency among CNO Stars, 401.
 Levato, H., *See Arnal et al.*
 Livi, S.H.B., *See Ducati et al.*
 Lizano, S. and F.H. Shu, Heating of Molecular Cloud Cores, 587.
 Loiseau, N., E. Hummel, R. Beck, and R. Wielebinski, Small Scale Structure of Polarized Emission in M31, 167.
 Loiseau, N., W. Reich, R. Wielebinski, and W. Münch, A Deep Radio Survey of the North Ecliptic Pole Region at 11 cm, 514.
 López, A., *See Laville et al.*
 López, D., *See Laville et al.*
 López, J.A., and E. de Lara, The Velocity Structure of the Bipolar Planetary Nebula 19W32, 559.
 López, R., *See Isern et al.*
 López Arroyo, M., *See Arellano Ferro et al.*
 López de Coca, P., *See González et al.*
 López de Coca, P., *See Rolland et al.*
- MacConnell, D.J., R.F. Wing, and E. Costa, CCD Spectroscopy of New M-Type Supergiants in the Southern Milky Way, 367.
 MacConnell, D.J., *See Wing et al.*
 Magris C., G. and G. Bruzual A., Internal Extinction in Spiral Galaxies. Inclination Dependence, 156.
 Malaroda, S., *See Levato et al.*
 Mampaso, A., P. Pişmiş, J.M. Vilchez, and J.P. Phillips, The Bipolar H II Region S201, 474.
 Mampaso, A., *See Carballo et al.*
 Mampaso, A., *See Manchado et al.*
 Mampaso, A., *Véase Vilchez y Mampaso.*
 Manchado, A., A. Mampaso, y S.R. Pottasch, Gradientes de abundancias en la nebulosa planetaria A78, 528.
 Marín, Z., *See Calvet and Marín.*
 Marques dos Santos, P., *See Epchtein et al.*
 Marraco, H.G., *See Clocchiatti and Marraco.*
 Martín, M.C., *Véase Bajaja et al.*
 Martín, M.C., *Véase Silva et al.*
 Martos, M.A., *See Allen and Martos.*
 Martos, M.A., *See Allen and Martos.*
 Martos, M.A., *See Allen et al.*
 Massey, P., *See Torres and Massey.*
 Matsuura, O.T., *See Epchtein et al.*
 Mattos, M.E., *See Clauzet and Mattos.*
 Maupomé, L., On the Astronomy in the Mesoamerican Calendar. Preliminary Remarks, 43.
 Maza J., y M.T. Ruiz, Nuevas galaxias Seyfert 1 australes, 101.
 Maza, J., *See Ruiz and Maza.*
 Mazzaro, J., *Véase Silva et al.*
- Mello, G.F.P., *See Quast et al.*
 Mello, G.F.P., *See Torres et al.*
 Mello, S.F., *Véase Balthazar et al.*
 Melnick, J., and M. Moles, On the Origin of the Dipole Anisotropy of the Cosmic Microwave Background: Beyond the Hydra-Centaurus Supercluster, 72.
 Melnick, J., R. Terlevich, and M. Moles, Giant H II Regions as Distance Indicators, 158.
 Melnick, J., *See Dottori et al.*
 Méndez, M., *Véase Forte et al.*
 Mendoza, B., *Véase Bravo et al.*
 Mendoza, C., *Véase Ferrín et al.*
 Mendoza, C., *Véase Naranjo et al.*
 Mendoza, E.E., *See Chavarría-K. et al.*
 Mendoza G., C.X., *See Mendoza and Mendoza G.*
 Mendoza T., E., Reactivación del radiointerferómetro solar de Tonantzintla, 759.
 Mendoza V., E.E. Photometric Classification of Early Type Stars with Hydrogen Emission Lines, 310.
 Mendoza V., E.E., A Preliminary Report on Photometric Measurements of the He I line λ 10830Å, 385.
 Mendoza V., E.E., and C.X. Mendoza G., The Féry Spectrograph Updated, 763.
 Menzies J., *See Solivella et al.*
 Michel, R., *See Alvarez et al.*
 Mirabel, I.F., *See Franco et al.*
 Mirabel, I.F., *Véase Silva et al.*
 Miranda, A.I., and M. Rosado, Nebulae Associated with Of Stars: S60, S108 and Of #23, 479.
 Moles, M., *See Melnick et al.*
 Moles, M., *See Melnick et al.*
 Morel, P.J., *Véase Chapellier et al.*
 Moreno C., M.A., Estudio cinemático de la región H II S155: algunos resultados, 488.
 Moreno, H., A. Gutiérrez-Moreno, C. Torres, and E. Wenderoth, Photographic and Spectroscopic Observations of Southern Planetary Nebulae, 520.
 Moreno, H., *See Gutiérrez-Moreno et al.*
 Morras, R., *Véase Bajaja et al.*
 Morras, R., *Véase Silva et al.*
 Morrell, N., *See Arnal et al.*
 Morrell, N., *See Levato et al.*
 Morrell, N., *See Levato et al.*
 Münch, W., *See Loiseau et al.*
 Muñoz Tuñón, C., J. Beckman, and M. Prieto, NGC 6946: Kinematics of the Starburst in the Circum-nuclear Zone, 144.
 Muñoz Tuñón, C., *See Acosta et al.*
 Muñoz Tuñón, C., *See Beckman et al.*
- Naranjo, O., F. Fuenmayor, I. Ferrín, P. Bulka, y C. Mendoza, Espectroscopía del Cometa Halley, 668.
 Naranjo, O., *Véase Ferrín et al.*
 Navarro, S., R. Costero, A. Serrano P.G., y L. Carrasco, Espectrofotometría de las estrellas simbióticas He2-417, He2-467 y He2-468, 339.
 Niemela, V.S., Binarias masivas en las Nubes de Magallanes, 390.
 Niemela, V.S., *See Gómez and Niemela.*
 Niemela, V.S., *See Cappa de Nicolau et al.*
- Ocegueda, J., *See Chavarría-K. et al.*
 Olalde, J.C., *Véase Silva et al.*
 Olano, C.A., *Véase Bajaja et al.*
 Olano, C.A., y W.G.L. Pöppel, Origen cinemático de las nubes oscuras de Tauro y de algunos cúmulos galácticos locales, 236.
 Ostriker, J.P. *See Aguilar et al.*
- Pani, A., *See Barral et al.*
 Paredes, J.M., Study of the Light Curve of LSI+61°303, 395.
 Parrao, L., *See Arellano Ferro et al.*
 Parrao, L., *See Rolland et al.*

- Pastoriza, M.G., *See* Bergmann and Pastoriza.
 Pastoriza, M.G., *See* Ducati *et al.*
 Payne, H., *See* Turner *et al.*
 Peimbert, M., Chemical Evolution of Galaxies, 166.
 Peimbert, M. and S. Torres-Peimbert, Chemical Composition of Type I Planetary Nebulae. Collisional Excitation Effects on the He I Line Intensity, 540.
 Peimbert, M., S. Torres-Peimbert, and J. Fierro, Physical Conditions of H II Regions in the Spiral Galaxy M101 and the Pre-galactic Helium Abundance, 165.
 Peimbert, M., N. Ukita, T. Hasegawa, and J. Jugaku, Radio Observations of M17 and the Orion Nebula: the He/H Ratio, 512.
 Peniche, R., and J.H. Peña, A Photometric Study of Short Period Variable Stars in Open Clusters, 420.
 Peniche, R., *See* Barral *et al.*
 Peniche, R., *See* Garrido *et al.*
 Peniche, R., *See* Peña *et al.*
 Peña, J.H., R. Peniche, S.F. González, and M.A. Hobart, Period Variations in SX Phe Stars: CY Aqr, DY Peg and HD 94033, 429.
 Peña, J.H., *See* Barral *et al.*
 Peña, J.H., *See* Garrido *et al.*
 Peña, J.H., *See* González *et al.*
 Peña, J.H., *See* Peniche and Peña.
 Peña, J.H., *See* Rolland *et al.*
 Peña, M., M.T. Ruiz, and M. Rubio, Spectrophotometric Observations of Very Low Ionization H II Regions in the LMC, 178.
 Peña, M., and S. Torres-Peimbert, JHK Photometry of Compact Planetary Nebulae, 534.
 Pérez-Enríquez, R., *Véase* Bravo *et al.*
 Pérez-Peraza, J., M. Alvarez-Madrigal, and A. Sánchez, Particle Acceleration in Cometary Processes of Magnetic Reconnection, 682.
 Pérez-Peraza, J., M. Alvarez-Madrigal, and F. Rivero, Numerical Analysis of the Azimuthal Transport of Solar Particles, 693.
 Pérez-Peraza, J. and A. Gallegos-Cruz, Solutions of the Fokker-Planck Equation for the Energy Distributions of Suprathermal Electrons, 705.
 Pérez-Peraza, J., *See* Gallegos and Pérez-Peraza.
 Pérez-Peraza, J., *See* Laville *et al.*
 Perrier, C., *See* Carrasco *et al.*
 Perrier, C., *See* Chelli *et al.*
 Phillips, J.P., *See* Mampaso *et al.*
 Picazzio, E., *See* Epchtein *et al.*
 Pimentel, P., *See* Recillas-Cruz *et al.*
 Pişmiş, P., La astronomía en México: su etapa moderna, 35.
 Pişmiş, P., Comments on Activity in Galactic Nuclei at all Scales of Energetics, 108.
 Pişmiş, P., and A.M. Cervantes, Statistical Properties of Lobes and Jets. I. In Galaxies with known Hubble Types, 168.
 Pişmiş, P., I. Hasse, and A. Quintero, The Emission Object Sharpless 207; is it a Planetary Nebula? 483.
 Pişmiş, P., *See* Mampaso *et al.*
 Pöppel, W.G.L., *Véase* Bajaja *et al.*
 Pöppel, W.G.L., *Véase* Silva *et al.*
 Pöppel, W.G.L., *Véase* Olano y Pöppel.
 Pottasch, S.R., *Véase* Manchado *et al.*
 Poveda, A., *See* Allen *et al.*
 Prieto, M., *See* Acosta *et al.*
 Prieto, M., *See* Beckman, *et al.*
 Prieto, M., *See* Muñoz Tuñón *et al.*
 Quast, G.R., C.A.P.C.O. Torres, R. de la Reza, and G.F.P. Mello, Isolated Post-T Tauri Stars, 360.
 Quast, G.R., *See* Torres *et al.*
 Quintero, A., *See* Barral *et al.*
 Quintero, A., *See* Pişmiş *et al.*
 Rabolli, M., Intercambio de cúmulos globulares en cúmulos de galaxias, 79.
 Ramos, M.P., *Véase* Cruz-González *et al.*
 Recillas-Cruz, E., L. Carrasco, A. Serrano P.G., and I. Cruz-González, Near Infrared JKH Photometry of Clusters of Galaxies. I. E, S0 and S Galaxies in the Coma Cluster, 77.
 Recillas-Cruz, E., P. Pimentel, and A. Serrano P.G., Environmental Variation of Radius with ρ_{Gal} in the Coma Cluster, 78.
 Recillas-Cruz, E., and A. Serrano P.G., On the Relation between Radius, Luminosity, Surface Brightness and Galaxian Surface Density in E galaxies, 170.
 Recillas-Cruz, E., *See* Cruz-González *et al.*
 Recillas-Cruz, E., *See* Serrano P.G. *et al.*
 Reich, W., *See* Loiseau *et al.*
 Ríos Berumen M., *See* Barral *et al.*
 Ríos Herrera M., *See* Barral *et al.*
 Rivero, F., *See* Pérez-Peraza *et al.*
 Rodríguez, L.F., *See* Curiel *et al.*
 Rodríguez, L.F., *See* Gómez *et al.*
 Rolland, A., P. López de Coca, M.A. Hobart, J.H. Peña, and L. Parrao, Photometric Study of the Delta Scuti Star Delta Serpentis, 419.
 Rolland, A., *See* González *et al.*
 Rosado, M., *See* Miranda and Rosado.
 Rossi, S.C.F., and P. Benevides-Soares, Quasi-Statical Solution of Early Homogeneous Nucleation Phases, 624.
 Roth, M., M. Tapia, and M.T. Ruiz, The Great Carina Nebula: Normal or Abnormal Extinction? 612.
 Roth, M., *See* Tapia *et al.*
 Rozyczka, M., *See* Franco *et al.*
 Rozyczka, M., *See* Tenorio-Tagle *et al.*
 Rubio, M., Molecular Clouds in the Large Magellanic Cloud, 188.
 Rubio, M., *See* Peña *et al.*
 Ruiz, E., *Véase* Angeles *et al.*
 Ruiz, M.T. and J. Maza, Discovery of Very Low Luminosity Stars, 381.
 Ruiz, M.T., *Véase* Maza y Ruiz.
 Ruiz, M.T., *See* Peña *et al.*
 Ruiz, M.T., *See* Roth *et al.*
 Ruiz, M.T., *See* Tapia *et al.*
 Ruiz Schneider, E., Instrumentación astronómica en México, 717.
 Sacristán, E., *Véase* Angeles *et al.*
 Sagnier, J.L., *Véase* Balthazar *et al.*
 Sahade, J., Discurso Inaugural, 15.
 Sahade, J., *See* Brandi *et al.*
 Sahade, J., *See* Solivella *et al.*
 Sala, F., Galactic Kinematics: an Axi-symmetric Time-dependent Model with Separable Potential, 195.
 Sala, F., *See* Cortés and Sala.
 Sánchez, A., *See* Pérez-Peraza *et al.*
 Santos, J.F., *See* Ducati *et al.*
 Sareyan, J.P., *See* Alvarez *et al.*
 Sareyan, J.P., *Véase* Chapellier *et al.*
 Sareyan, J.P., *Véase* González *et al.*
 Sarmiento, G., A., *See* Serrano P.G. *et al.*
 Sasselov, D.D., *See* Arellano Ferro *et al.*
 Sato, M., and T. Yokoyama, A First Order Analytical Elliptic planar Model for 3/1 Resonance Problem, 641.
 Sato, M., *See* Yokoyama *et al.*
 Scalise, Jr. E., *Véase* Abraham *et al.*
 Scalise, Jr. E., *Véase* Sestokas Filho and Scalise.
 Schmidt, A., *See* Ducati *et al.*
 Schmidt, A.A., C.A.Z. Vasconcellos, and H.A. Dottori, Cosmological Mass Density and the Regge Law, 66.
 Schmidt, A.A., and H.A. Dottori, Nuclear Stellar Population of NGC 772 (Arp 78), 149.
 Schroder, M.F.S., *See* Ducati *et al.*
 Serrano P.G., A., A. González, J.F. Barral, E. Recillas-Cruz, and A. Sarmiento G., Chaotic Regions and Invariant Manifolds in Models for Elliptical Galaxies, 80.

- Serrano P.G., A., *Véase* Navarro *et al.*
 Serrano P.G., A., *See* Recillas-Cruz and Serrano P.G.
 Serrano P.G., A., *See* Recillas-Cruz *et al.*
 Serrano P.G., A., *See* Recillas-Cruz *et al.*
 Sessin, W., Aspects of the Phase-Portrait of Resonant Problems, 631.
 Sessin, W., *See* Bevilacqua and Sessin.
 Sestokas Filho, B., e E. Scalise Jr., Variabilidade temporal de fontes maser de vapor d'água associadas às regiões H II, 513.
 Shapiro, P.R., and H. Kang, Hydrogen Molecules and the Radiative Cooling of Pregalactic Shocks II: Low Velocity Shocks at High Redshift, 58.
 Shu, F.H., *See* Lizano and Shu.
 Silva, A.M., E. Bajaja, R. Morras, J.C. Cersosimo, M.C. Martín, E.M. Arnal, W.G.L. Pöppel, F.R. Colomb, J. Mazzaro, J.C. Olalde, V. Boriakoff e I.F. Mirabel, Radio observaciones del OH en la coma del Cometa Halley desde el hemisferio sur, 670.
 Simonneau, E., *See* Isern *et al.*
 Sisteró, R.F., *Véase* Clariá *et al.*
 Solivella, G.R., J. Menzies, Y. Kondo, and J. Sahade, Spectroscopic Study of the Binary System HD 153919, 417.
 Stalio, R., *See* Carrasco *et al.*
 Steigman, G., In Search of Primordial Helium, 71.
 Steiner, J.E. e S.D. Kirhakos, Inclinação e obscurecimento em galáxias de Seyfert, 103.
 Steiner, J.E., D. Cieslinski, and F.J. Jablonski, Looking for New Cataclysmic Variables, 436.
 Stencel, R.E., *Véase* Hagen *et al.*
 Stock, J., y C. Abad, Análisis del campo de distorsión del catálogo astrográfico, 448.
 Tapia, M., M. Roth, L. Carrasco, and M.T. Ruiz, On the Infrared Emission of the Exciting Star of the Herbig-Haro Objects 1 and 2, 517.
 Tapia M., *See* Roth *et al.*
 Tateyama, C.E., *Vea-se* Abraham *et al.*
 Tejada de Vargas, C., *Véase* Cobos y Tejada de Vargas.
 Tenorio-Tagle, G., P. Bodenheimer, M. Rozyczka, and J. Franco, Collisions of High-Velocity Clouds with a Galactic Disk, 241.
 Tenorio-Tagle, G., *See* Franco *et al.*
 Terlevich, R., *See* Melnick *et al.*
 Terzian Y., *See* Turner *et al.*
 Tinoco, S. y S. Cuevas, Diseño de un interferómetro de rotación de frente de onda para obtención de imágenes de alta resolución espacial, 756.
 Torra, J., *See* Gómez and Torra.
 Torres, A.V., and P. Massey, An Atlas of Optical Spectrophotometry of Southern Wolf-Rayet Carbon Stars, 287.
 Torres C., *See* Moreno *et al.*
 Torres, C.A.P.C.O., *See* Quast *et al.*
 Torres, C.A.P.C.O., G.R. Quast, R. de la Reza, and G.F.P. Mello, V4046 Sgr—An Isolated Binary Post-T Tauri Star, 361.
 Torres-Peimbert, S., *See* Peimbert and Torres-Peimbert.
 Torres-Peimbert, S., *See* Peimbert *et al.*
 Torres-Peimbert, S., *See* Peña and Torres-Peimbert.
 Turner, K.C., Y. Terzian, and H. Payne, Galactic Confusion and the Detection of 1665 and 1667 MHz OH Lines in the Neighborhood of Planetary Nebulae, 567.
 Ukita, N., *See* Peimbert *et al.*
 Valdés-Galicia J., *Véase* Bravo *et al.*
 Valtier, J.C., *Véase* Chapellier *et al.*
 Vasconcellos, C.A.Z., *See* Schmidt *et al.*
 Vega, E.I., *Véase* Forte *et al.*
 Veiga, C.H., *Vea-se* Vieira-Martins e Veiga.
 Veiga, C.H., *See* Vieira-Martins *et al.*
 Viegas-Aldrovandi, S.M. and M. Contini, Active Galactic Nuclei: Narrow Emission Line Profiles, 83.
 Viegas-Aldrovandi, S.M. and R.B. Gruenwald, The effect or Relativistic Electrons on the Narrow Emission Lines of AGN, 87.
 Viegas-Aldrovandi, S.M., *See* Andrade *et al.*
 Viegas-Aldrovandi, S.M., *Vea-se* Andrade *et al.*
 Viegas-Aldrovandi, S.M., *See* Laguna and Viegas-Aldrovandi.
 Vieira-Martins, R. e C.H. Veiga, Determinação astrométrica de posições com filmes do "FSO-B Survey", 456.
 Vieira-Martins, R., C.H. Veiga, and D. Lazzaro, Observations of the Satellites of Uranus, 642.
 Vilas Boas, J.W.S., *Vea-se* Abraham *et al.*
 Vilchez, J.M., y A. Mampaso, La nebulosa cometaria 1548C27 515.
 Vilchez, J.M., *See* Mampaso *et al.*
 Vogt, N., *See* Barrera and Vogt.
 Wampler, J., *See* Barral *et al.*
 Welch, W.J., *See* Carral *et al.*
 Wenderoth, E., *See* Moreno *et al.*
 Wielebinski, R., *See* Loiseau *et al.*
 Wielebinski, R., *See* Loiseau *et al.*
 Wing, R.F., D.J. MacConnell, and E. Costa, New M-Type Supergiants in the Southern Milky Way, 362.
 Wing, R.F., *See* MacConnell *et al.*
 Wright, M.C.H., *See* Carral *et al.*
 Yokoyama, T., M. Sato, J. Koiller, and J.M. Balthazar, Some instabilities in the 3/1 Resonant Problem, 636.
 Yokoyama, T., *Vea-se* Balthazar *et al.*
 Yokoyama, T., *See* Sato and Yokoyama.
 Zepka, A.F. and H.A. Dottori, Morphology of LMC Clusters, 172.
 Zinnecker, H., *See* Carrasco *et al.*