

$\log \text{He}/\text{H} = 10.96 \pm 0.05$, $\log \text{O}/\text{H} = 8.10 \pm 0.10$, $\log \text{Ne}/\text{H} = 7.5 \pm 0.2$ and $\log \text{Ar}/\text{H} = 5.8 \pm 0.2$. The O, Ne and Ar appear underabundant by a factor of 5, relative to solar abundances. These results indicate that PN 6 -41.1 is a population II object with chemical composition comparable to the SMC H II regions and very similar to the only known C poor halo planetary nebula DDDM-1.

From nebular parameters we have estimated that the distance to the object is in the range $2.8 \text{ kpc} \leq d \leq 18 \text{ kpc}$, corresponding to a galactic plane distance range of $1.9 \text{ kpc} \leq |z| \leq 12 \text{ kpc}$. On the other hand, from the characteristics of the central star, we derived a distance $d \cong 7.7 \text{ kpc}$, with $|z| \cong 5.1 \text{ kpc}$, which we consider is the most probable distance to the object. This distance estimate places the nebula in the galactic halo. The complete version of this work appears in *Revista Mexicana Astron. Astrof.*, 17, 1989.

AGE STRUCTURE OF REFRACTORY INTER-STELLAR DUST AND ISOTOPIC CONSEQUENCES

Donald D. Clayton and Paul A. Scowen

Department of Space Physics and Astronomy
Rice University

Kurt Liffman

NASA Ames Research Center

We compute the mean age of dust particles using a sputtering and recycling Monte Carlo model developed by Liffman, and Clayton, (1988, *Proc. Lunar Planet. Sci. Conf.*, 18, 637). Each particle mean age is defined not as the time it has existed, but rather as the mass-weighted existence times of its parts (core plus shell) at $t = 6 \text{ Gyr}$ when the Solar System formed in our models. We show that galactic evolution generates a correlation between particle size and mean age. This is a mean correlation, applying to large numbers of particles binned according to size rather than to individual particles, whose mean ages fluctuate statistically. The cosmochemical consequence is that if interstellar particles can be dynamically sorted into separate size populations during the aggregation history of solar system bodies, the collections of larger grains will constitute matter that is chemically older than collections of smaller grains. This macroscopic age difference generates isotopic anomalies by virtue of the time dependence of the secondary/primary nucleosynthesis yields (Clayton, 1988, *Ap. J.*, 334, 191). The most important example is that an aggregate of refractory oxides is several percent richer in ^{16}O than is the solar gas, ranging up to 10% richer in Al_2O_3 if Al is also concentrated into larger-than-average particles. This history may explain the 5% richness of ^{16}O within meteoritic Al-rich inclusions. We compare our results with three different prescriptions for the sputtering of interstellar dust.

IUE OBSERVATIONS OF WOLF-RAYET BINARIES IN THE GALAXY, THE LMC AND THE SMC

G. Koenigsberger

Instituto de Astronomía
Universidad Nacional Autónoma de México

L.H. Auer

Los Alamos National Laboratory

A.F.J. Moffat

Dept. de Physique, Université de Montréal, Canada

Due to the large extent of the winds of W-R stars, atmospheric eclipses are evident at *UV* line frequencies in binary systems containing an O-star companion. In 8 of the galactic WR's thus far studied with the *IUE* satellite, the presence of very numerous, closely spaced lines of Fe IV, Fe V and Fe VI lead to pseudo-continua which are most evident during the atmospheric eclipses, even in the lowest inclination binary systems. We report on the results of *IUE* observations of the systems HD 36402 in the LMC, and HD 5980 and Sk 188 in the SMC where no evidence is found for the iron pseudo-continua, although atmospheric eclipse effects are relatively strong at line frequencies corresponding to ions of N IV, C IV and O IV. This is consistent with the lower heavy-metal abundances in the Clouds with respect to the Solar vicinity.

THE DISTANCES TO HIGH VELOCITY CLOUDS: A MODEL FOR THEIR CONFINEMENT

Julia Espresate, José Franco, and Jorge Cantó

Instituto de Astronomía
Universidad Nacional Autónoma de México

The assumption that the High Velocity Clouds are embedded in the hot galactic gaseous halo, and in pressure equilibrium with it, permits the determination of a distance to the clouds.

Being in pressure equilibrium with the surrounding medium means that the total external pressure equals the pressure within the cloud. That is, we are assuming no expansion or contraction of the cloud.

In this model the external pressure has two components. One is the thermal pressure of the hot surrounding gas, and the other is the ram pressure due to the drag of the material they find in their trajectories. This external pressure depends on the distance because the distribution of the coronal gas is not uniform.

From the parameters observed for each individual cloud (column density, line width, angular size and velocity with respect to the galactic standard of rest) we obtained the internal pressure as a function of distance