

EVIDENCE FOR THE PRESENCE OF A WARPING OF THE IONIZED GAS LAYER DERIVED FROM H166 α EMISSION OBSERVATIONS IN THE OUTER GALAXY

I.N. Azcárate¹, J.C. Cersosimo¹, and F.R. Colomb¹

Received 1989 November 8

RESUMEN

Se presentan resultados de un relevamiento de la emisión de la línea H166 α en el rango de longitudes galácticas $270^\circ \leq l \leq 300^\circ$ y para tres latitudes galácticas $b = 0.0^\circ$, $b = \pm 0.5^\circ$. De los mismos, resulta que hay evidencia de un alabeo de la capa de gas ionizado en el plano galáctico, en las partes exteriores de la Galaxia.

ABSTRACT

Results from an H166 α emission survey in the Galactic longitude range $270^\circ \leq l \leq 300^\circ$ and for three Galactic latitudes ($b = 0.0^\circ$, $b = \pm 0.5^\circ$), are presented. From these results, there is evidence for the presence of a warping of the ionized gas layer in the galactic plane, in the outer Galaxy.

Key words: GALAXY-STRUCTURE – RADIO LINES-RECOMBINATION

I. INTRODUCTION

The existence of a warping of the H I layer in the outer parts of the Milky Way, has been known since several years ago, from early 21-cm surveys (Burke 1957; Kerr 1957; Westerhout 1957; Oort, van der Kruit, and Westerhout 1958). The atomic hydrogen layer is systematically warped from the galactic equator near $b = 0.0^\circ$.

On the other hand, these systematic deviations in a flat disk are a common characteristic of the H I morphology in the outer regions of spiral galaxies, as has become clear from several works; for example the nearest large spirals H31 and H33 are both warped (Brinks and Burton 1984). Kerr *et al.* (1986) used the Parkes 18m-telescope to effectively extend the coverage provided by the Weaver and Williams (1973) H I survey in the range $-10^\circ \leq b \leq 10^\circ$ in the north to include all galactic longitudes. Henderson, Jackson, and Kerr (1982) used the combined $|b| < 10^\circ$ material to obtain a global description of the detailed shape of the warped H I layer. This general warping above the galactic plane in the north and below the plane in the south is clearly shown in the paper by Henderson *et al.* (1982).

From CO ($J = 1 \rightarrow 0$) observations in the range $270^\circ \leq l \leq 300^\circ$ and latitudes $-5^\circ \leq b \leq 5^\circ$, Grabelsky *et al.* (1987) show that there is a similar warping of the molecular layer in the galactic plane in the outer Galaxy. Between galactocentric distances $R = 10.5$ kpc and $R = 12.5$ kpc, the CO midplane dips from $z = -46$ to -167 pc below the galactic plane. Clemens, Sanders, and Scoville (1988), have shown something similar, from CO observations in the first quadrant (warping of the CO layer away from $b = 0.0^\circ$).


In the present paper, we give some results of an H166 α emission survey in the galactic coordinate range $270^\circ \leq l \leq 300^\circ$, $b = 0.0^\circ$, $b = \pm 0.5^\circ$. This survey was carried out, on the one hand, to complete the published H166 α emission survey for $300^\circ \leq l \leq 360^\circ$, $b = 0.0^\circ$ (Cersosimo *et al.* 1989) and, on the other hand, with the main purpose of searching for evidence of a similar warping of the low-density ionized gas layer in the outer Galaxy, away from $b = 0.0^\circ$. We will describe below the observations carried out with this purpose in the Instituto Argentino de Radioastronomía.

II. THE OBSERVATIONS

H166 α line emission has been observed in the galactic longitude range $270^\circ \leq l \leq 300^\circ$, in steps at one degree in longitude and for three galactic latitudes: $b = 0.0^\circ$, $b = \pm 0.5^\circ$ for each longitude.

The observations were made with the 30m-

1. Members of the Carrera del Investigador Científico Tecnológico del Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.



diameter antenna of the Instituto Argentino Radioastronomía, with Gaussian beam of $34'$ 21-cm. The aperture efficiency was 0.6 and beam efficiency was 0.87 (Loiseau 1979). The filter bank spectrometer had 84 filters of widths 75 kHz giving a velocity resolution of 15.8 km s^{-1} . The system temperature on cold sky was 84 K. The local oscillator was switched in frequency by an amount less than the observing bandwidth, in order to obtain two independent spectra, which were then averaged.

The profiles had an integration time of 4-6 minutes. That represented a very long observation time for the complete survey. The 'r.m.s.' noise of the profiles was $\approx 7\text{-}8 \text{ mK}$, with a minimum detectable temperature of 20 mK. The instrumental baselines were removed by using third, fourth or fifth order polynomials.

III. RESULTS

Figure 1 shows some of the observed profiles







Fig. 1. Some profiles obtained from H166 α observations in the range $270^\circ \leq l \leq 300^\circ$ and three galactic latitudes, ($b = 0.0^\circ, \pm 0.5^\circ$). In two of the points shown there is no signal detected.

in all the observed profiles, the result is that the H166 α emission was detected clearly in relatively small positions: 4 for $b = 0.0^\circ$, 4 for $b = -0.5^\circ$, 1 for $b = -0.5^\circ$, (for $280^\circ \leq l \leq 300^\circ$ and the three galactic latitudes). No detection was positive, above the 3σ level, for the range $270^\circ \leq l \leq 280^\circ$. The peak temperature of the obtained profiles, together with their galactic coordinates are shown in Table 1. The range $270^\circ \leq l \leq 279^\circ$ is not considered in the Table 1, since there was no positive detection. As an immediate result, we almost had no detections for $b = +0.5^\circ$. That is in good agreement with the previously known warping of different gaseous layers (H I, CO, H₂) in the outer galactic plane, away from $b = 0.0^\circ$ (below the plane in the south and above in the north). That phenomenon could be also present for the low-density ionized gas, according to our H166 α observations.

IV. COMMENTS

In this paper, we have shown evidence of the presence of the gaseous layer warping in the Milky Way, also for ionized gas (at least for the low density component). It is known that warping is generally also present in the outer regions of spiral galaxies. These systematic deviations from flatness seem to be a common aspect of spiral galaxy morphology, and their presence in the Milky Way, is established beyond doubt. We think that the present paper is an observational contribution to confirm this characteristic, in relation to the ionized gas, common to several spiral galaxies.

REFERENCES

Brinks, E. and Burton, W.B. 1984, *Astr. and Ap.*, **141**, 195.

TABLE 1

SOME OBSERVED PARAMETERS IN THE H166 α OBSERVATIONS

Galactic Longitude $l(^{\circ})$	Galactic Latitude $b(^{\circ})$	Peak Line Temperature T_A (mK)	Galactic Longitude $l(^{\circ})$	Galactic Latitude $b(^{\circ})$	Peak Line Temperature T_A (mK)
280	-0.5	< 20.	291	0.0	< 20.
281	-0.5	< 20.	292	0.0	< 20.
282	-0.5	< 20.	293	0.0	< 20.
283	-0.5	< 20.	294	0.0	< 20.
284	-0.5	284.	295	0.0	< 20.
285	-0.5	< 20.	296	0.0	< 20.
286	-0.5	31.	297	0.0	< 20.
287	-0.5	167.	298	0.0	< 20.
288	-0.5	118.	299	0.0	< 20.
289	-0.5	< 20.	300	0.0	< 20.
290	-0.5	< 20.	280	0.5	< 20.
291	-0.5	< 20.	281	0.5	< 20.
292	-0.5	< 20.	282	0.5	< 20.
293	-0.5	< 20.	283	0.5	< 20.
294	-0.5	< 20.	284	0.5	< 20.
295	-0.5	< 20.	285	0.5	< 20.
296	-0.5	< 20.	286	0.5	< 20.
297	-0.5	< 20.	287	0.5	< 20.
298	-0.5	< 20.	288	0.5	< 20.
299	-0.5	< 20.	289	0.5	< 20.
300	-0.5	< 20.	290	0.5	< 20.
280	0.0	23.	291	0.5	< 20.
281	0.0	< 20.	292	0.5	< 20.
282	0.0	< 20.	293	0.5	22.
283	0.0	< 20.	294	0.5	62.
284	0.0	82.	295	0.5	< 20.
285	0.0	< 20.	296	0.5	< 20.
286	0.0	< 20.	297	0.5	< 20.
287	0.0	44.	298	0.5	< 20.
288	0.0	30.	299	0.5	< 20.
289	0.0	< 20.	300	0.5	< 20.
290	0.0	< 20.			

- Burke, B.F. 1957, *A.J.*, **62**, 90.
Cersosimo, J.C., Azcárate, I.N., Hart, L., and Colomb, F.R. 1989, *Astr. and Ap.*, **208**, 239.
Clemens, D.P., Sanders, D.B., and Scoville, N.Z. 1988, *Ap. J.*, **327**, 139.
Grabelsky, D.A., Cohen, R.S., Bronfman, L., Thaddeus, P., and Hay, J. 1987, *Ap. J.*, **315**, 122.
Henderson, A.P., Jackson, P.D., and Kerr, F.J. 1982, *Ap. J.*, **283**, 116.
Kerr, F.J. 1957, *A.J.*, **62**, 93.
Kerr, F.J., Bowers, P.F., Jackson, P.D., and Kerr, M. 198, *Astr. and Ap. Suppl.*, **66**, 373.
Loiseau, N. 1979, *Informe Técnico del IAR*, No. 12.
Oort, J.H. Kerr, F.J., and Westerhout, G. 1958, *M.N.R.A* **118**, 379.
Weaver, H.F. and Williams, D.R.W. 1973, *Astr. and A Suppl.*, **8**, 1.
Westerhout, G. 1957, *Bull. Astr. Inst. Netherlands*, **13**, 20

I.N. Azcárate, J.C. Cersosimo, and F.R. Colomb: Instituto Argentino de Radioastronomía, Casilla de Correos No. 5, (1894) Villa Elisa, Prov. de Buenos Aires, Argentina.