

ON WOLF-RAYET STARS, BLACK HOLES AND CO-ROTATING BINARY SYSTEMS

A. Mosqueda¹ and G. Koenigsberger¹

Instituto de Astronomía
Universidad Nacional Autónoma de México

Received May 3 1990

RESUMEN

Se analiza el problema de la formación de agujeros negros en sistemas binarios que corrotan. Se utilizan el valor crítico de a/m de Miller y De Felice y la hipótesis de corrotación para hallar el periodo orbital crítico de un sistema binario con una estrella Wolf-Rayet, justo antes del colapso de ésta. Se comparan estos periodos críticos con datos observacionales para identificar aquellos sistemas WR que puedan engendrar agujeros negros en el futuro.

ABSTRACT

The problem of black hole formation in a co-rotating binary system is analyzed. The Miller and De Felice critical value of a/m and the co-rotation hypothesis, are used to find critical orbital periods in a Wolf-Rayet system just prior to the collapse of the more evolved star. These critical periods are compared with observational data to find WR binary systems that may produce black holes in the future.

Key words: **BLACK HOLES – STARS-BINARIES – STARS-WOLF-RAYET**

I. INTRODUCTION

The Wolf-Rayet (WR) stars in massive binary systems are presumed to be progenitors of massive X-ray binary systems, or systems containing collapsed companions (Van den Heuvel 1976). Since some of the masses derived for WR members of binary systems are very large, these stars are good candidates for progenitors of black holes (BH's). Thus, in addition to the interest one may have in the WR phenomenon itself, an understanding of these stars has relevance to the problem of the existence of BH's in massive binary systems.

The general properties and evolutionary scenarios for the formation of WR stars are described in Abbott and Conti (1987) and Chiosi and Maeder (1986), according to which the massive WR stars evolve from O-stars more massive than about $40 M_{\odot}$, and consist of He-burning cores with severely altered surface chemical abundances: significant underabundance of H, overabundance of He, and N or He and C, relative to solar abundances. As massive, post-H-main sequence objects their expected lifetimes do not exceed a few hundred thousand

years, after which they produce a supernova explosion.

Black holes are believed to result from the collapse of a massive star, following a supernova explosion. However, there are certain conditions that must be satisfied for the collapse to lead to a BH; in particular, the value of the specific angular momentum at the moment of collapse must be smaller than unity (Abramowicz and Lasota 1980). In itself, this condition does not seem to be a strong restriction for the formation of a BH. However, when one adds to this the generally adopted condition of co-rotation in close binary systems, constraints can be derived that restrict the number of close, massive binary systems in which a BH is allowed to form. In this paper we show that if we assume close binary systems to be co-rotating, and if no more than 40 % of the specific angular momentum of the exploding star is lost (Miller and De Felice 1985), then BH's can form only from binary WR systems with pre-explosion orbital periods larger than 15 days. This would have important consequences on the interpretation of, for example, Cyg X-1, which contains the strongest BH candidate, and where the pre-explosion orbital period has been calculated to be 4.8 days (Sutantyo 1974).

1. Members of the Programa Universitario de Investigación y Desarrollo Espacial, UNAM.

II. CONSTRAINT ON THE ORBITAL PERIOD IMPOSED BY CO-ROTATION

A corollary of the third law of thermodynamics of black holes (Abramowicz and Lasota 1980) states that the ratio of specific angular momentum to gravitational mass must be less than unity in order for the black hole to form: i.e.,

$$\frac{a}{m} < 1 \quad , \quad (1)$$

with

$$a = \frac{l}{cm} \quad m = \frac{Gm}{c^2} \quad ;$$

where

l = angular momentum, c = speed of light, m = rest mass of the BH, and G = gravitational constant.

This restriction applies to the stellar remnant undergoing final collapse following the supernova explosion; i.e., once the ejecta has carried away a fraction of the specific angular momentum and mass. If the assumption is made that the amount of specific angular momentum and mass removed from the immediate progenitor is not constrained, then the condition $a/m < 1$ can always be achieved. However, Miller and De Felice, (MD, 1985) have shown that the pre-supernova value of a/m cannot be reduced by a factor larger than 2.5. This means that, in order for the collapse to lead to a black hole, the following condition on the immediate progenitor star (just prior to the supernova event) must be satisfied:

$$\frac{cl}{Gm^2} < 2.5 \quad . \quad (2)$$

This restriction applies under the following assumptions (MD):

1. The gravitational collapse is treated as that of a rotating fluid with axial symmetry.
2. The stellar magnetic field, pressure gradients, and energy losses by gravitational radiation are not taken into account.
3. The star is a rigid rotator.
4. Redistribution of angular momentum does not occur during collapse.

Using $l = I\omega$, $\omega = 2\pi/T$, one finds a critical rotation period for the pre-supernova star from (2):

$$T_c > \frac{2\pi Ic}{2.5Gm^2} \quad (3)$$

Thus, given the MD assumptions, a star rotating with $T < T_c$ just prior to the supernova explosion cannot engender a black hole.

If we consider a binary system consisting of two massive ($M_1, M_2 > 40 M_\odot$) stars, the hypothesis of co-rotation is represented by:

$$\frac{\Omega}{\omega} = 1 \quad , \quad (4)$$

where Ω is the orbital period of the system, and ω is the rotation period of the less massive of the two stars. If we now identify T_c with the orbital period condition 3 can be used to obtain a lower bound on the orbital periods of potential progenitors of binary systems containing black holes:

$$T_{orb} = T_c > \frac{2\pi Ic}{2.5Gm^2} \quad (5)$$

III. APPLICATION TO WR BINARY SYSTEMS

Let us consider a binary system consisting of a massive star which has not evolved far beyond the main sequence, and a less massive star which has reached its final evolutionary stages, and which we will identify as a Wolf-Rayet star. The latter is assumed to be a star consisting mostly of He.

The moment of inertia, I , in (5) is defined as

$$I = \int \rho r^2 dv \quad ;$$

where $\rho(r)$ is the interior density structure of the He star. Three different density profiles were adopted: a constant density, one that decreases linearly from the center to the surface, and one which decreases exponentially. These density profiles were adopted so as to be consistent with the assumptions of MI (i.e., rigid rotation) and to illustrate the dependence of T_c on the interior density distribution. The most realistic of these $\rho(r)$ distributions is the exponential one, according to the model of the $8 M_\odot$ core of a $22 M_\odot$ star just prior to final collapse illustrated by Schramm (Figure 2 in Iben, Renzini and Schramm 1977). In column 2 of Table 1 we list the moments of inertia derived for each of these density profiles. In column 3 we present the corresponding expression for T_c (in units of days). The values of T_c , as a function of the mass are plotted in Figure 1a (the same curves are replotted in Figures 1b-1d), where we have used the masses and radii of model He star given by Divine (1965) and Langer (1989). These curves represent the smallest orbital period a binary system can have, for a given mass of the WR star and where the post-supernova stellar remnant can collapse into a black hole. Since the mass of the

TABLE 1
CRITICAL PERIODS FOR THE DIFFERENT INTERIOR
DENSITY DISTRIBUTIONS

Density	M.Inertia	T_c (days)
$\rho = \rho_0 = \text{const.}$	$\frac{2}{5} M R_*$	$T_c = 12.76 (R/R_\odot)^2 (M/M_\odot)^{-1}$
$\rho = \rho_0 (1 - r/R_*)$	$\frac{8}{90} \pi \rho_0 R_*^5$	$T_c = 63.36 (R/R_\odot)^2 (M/M_\odot)^{-1}$
$\rho = \rho_0^{-\frac{r}{R_*}}$	$\frac{9}{100} \pi \rho_0 R_*^5$	$T_c = 260.36 (R/R_\odot)^2 (M/M_\odot)^{-1}$

TABLE 2
POSSIBLE BH PROGENITOR SYSTEMS ASSUMING
AN EXPONENTIAL INTERIOR DENSITY DISTRIBUTION

Number	Name	Mass WR	Ref.	Period (days)
1	CVSer	13–15 M_\odot	1,2	29.71
2	γ^2 Vel	18–19 M_\odot	1,2	78.50
3	HD 1909	18 15 M_\odot	1	112.80

1) Schulte-Ladbeck (1989); 2) Smith and Maeder (1989).

jecta of Type II supernovae has been estimated to be 5 M_\odot , (Chevalier 1977), this means that the possible black hole progenitors must have masses larger than 8 M_\odot given an upper limit of 3 M_\odot to the mass of neutron stars (White 1989). This means that only in binary systems with $T_c > 15$ days will form a black hole, adopting the exponential interior density distribution.

The observational data for WR stars are also plotted in Figures 1a-1d, each figure containing the data presented by different authors (1a, De Greve and De Loore 1975; 1b, Sutantyo and Dermawan 1981; 1c, Schulte-Ladbeck 1989; and 1d, Smith and Maeder 1988). The dotted line at 8 M_\odot is the above mentioned lower limit for the BH progenitor mass. The curve corresponding to an exponential interior density distribution provides the strongest constraint on the possible progenitors of BH's, and in Table 2 we list the three WR systems which satisfy these conditions. If one adopts a constant interior density distribution, the number of possible BH progenitors increases significantly; these are listed in Table 3 according to the data of Smith and Maeder (1989). However, as pointed out above, this density distribution is not realistic.

Also plotted in Figures 1a-1d is the position of the progenitor of Cyg X-1 as derived by the calculation of Sutantyo (1974), and which is in strong contradiction with the fact that this binary

TABLE 3
POSSIBLE BH PROGENITOR SYSTEMS ASSUMING
A CONSTANT INTERIOR DENSITY DISTRIBUTION

Number (WR)	Name (HD)	Mass WR (M_\odot)	Period (days)
9	63099	10	14.70
11	68273	19	78.50
21	90657	12	8.26
42	97152	11	78.50
47	E 311884	43	6.34
79	152270	7	8.89
127	186943	16	9.56
133	190918	15	112.80
148	197406	60	4.32
153	211853	14	6.69

system is the one most likely to contain a black hole, assuming an exponential interior density distribution.

IV. DISCUSSION AND CONCLUSIONS

Given two assumptions: 1) The maximum value of a/m for the progenitor of a black hole cannot exceed 2.5 and 2) the progenitor binary system is co-rotating, we find that only three WR binary systems




Fig. 1. Curves of critical T_c as a function of the progenitor's mass (i.e., the mass just prior to the final collapse and SN explosion), with the positions of the WR stars according to data of a) De Greve and De Loore 1975; b) Sutantyo and Dermawan 1981; c) Schulte-Ladbeck 1989; d) Smith and Maeder 1989. A * indicates the position of the progenitor of Cyg X-1 (Sutantyo 1974). This figure has been adapted from Mosqueda (1990).

have orbital periods which satisfy the constraint for the formation of a black hole. However, these same constraints lead to a major contradiction when applied to Cyg X-1.

The calculation of the predicted T_c vs mass shown in Figures 1 is based on these two assumptions, the weaknesses of which we now point out.

The first assumption is that the MD results for the maximum amount of specific angular momentum lost by the system are realistic. This leads to the adoption of the assumptions used by MD (listed in §II) for our analysis, and in particular, that of rigid rotation. However, if the pre-supernova stellar object is undergoing differential rotation, the redistribution of specific angular momentum during collapse becomes more probable. Although a reanalysis of the MD assumptions and analyses goes beyond the scope of this paper, we find that increasing the critical value of a/m to 12.5 (rather than 2.5) prior to collapse "allows" Cyg X-1 to contain a black hole and most of the WR

binaries to form one regardless of the interior density distribution adopted. Indeed, MD note that differential rotation probably would tend to increase the critical a/m and gravitational radiation could decrease this value.

The second assumption is that the binary system is co-rotating prior to the supernova explosion. The hypothesis of co-rotation in close binary systems is based on the work of Zahn (1966a,b,c.; 1975), van den Heuvel (1967), Lea and Margon (1973), and Sutantyo (1974), and applies to stars in or close to the main sequence. The phenomena associated with the evolution of a main sequence star towards a WR star are not yet well understood. In particular, the mechanisms by which an extraordinary fraction of the progenitor's mass is removed in order to expose inner, nuclear processed layers have not been studied in detail (the mass loss by stellar winds, as currently observed, cannot account for the removal of 70 – 80% of a progenitor star's mass). Is it not clear whether the resulting WR star may be

rotating with a period shorter than or larger than the orbital period. As pointed out by an anonymous referee, as the helium core contracts from a less dense progenitor that may have been in co-rotation, its rotation period will tend to speed up. However, as the WR star evolves, the presence of magnetic fields (Maheswaran and Cassinelli 1988) in addition to tidal interaction can lead to an increase of the orbital period. Furthermore, the probable episodic large mass loss events leading to the final stages of the WR phase can carry away angular momentum, slowing it down even more.

We thus conclude that before one can attempt to predict which WR systems can engender a black hole, one must have a better understanding of the evolution of close binary systems, and especially those aspects concerning the way in which the specific angular momentum of the WR component evolves with time.

We thank an anonymous referee for very useful comments and suggestions.

REFERENCES

- Abbott, D.C. and Conti, P.S. 1987, *Ann. Rev. Astr. and Ap.*, **25**, 113.
- Abramowicz, M.A. and Lasota, J.P. 1980, *Acta Astr.*, **30**, 35.
- Chevalier, R.A. 1977, *Ann. Rev. Astr. and Ap.*, **15**, 175.
- Chiosi, C. and Meader, A. 1986, *Ann. Rev. Astr. and Ap.*, **24**, 329.
- De Greve, J.P. and De Loore, C. 1975, *Astr. and Ap.*, **42**, 91.
- Divine, N. 1965, *Ap. J.*, **142**, 824.
- Iben, I., Renzini, A., and Scharmm, D.N. 1977, *Advanced Stages in Stellar Evolution*, Seventh Advanced Course of the Swiss Society of Astronomy and Astrophysics, Geneva Observatory Publ, p. 296.
- Langer, N. 1989, *Astr. and Ap.*, **210**, 93.
- Lea, S.M. and Margon, B. 1973, *Ap. (Letters)* **13**, 33.
- Maheswaran, M. and Cassinelli, J.P. 1988, *Ap. J.*, **335**, 931.
- Mosqueda, A. 1990, Tesis de Licenciatura, Fac. de Ciencias, UNAM, México.
- Miller, J.C. and De Felice, F. 1985, *Ap. J.*, **298**, 474.
- Schulte-Ladbeck, R.E. 1989, *A.J.*, **97**, 1471.
- Smith, F. and Maeder, A. 1989, *Astr. and Ap.*, **211**, 71.
- Sutantyo, W. 1974, *Astr. and Ap.*, **35**, 251.
- Sutantyo, W. and Dermawan, L. 1981, Second IAU Asia and Pacific Regional Meeting, Bandung.
- Van den Heuvel, E.P.J. 1967, *Bull. Astr. Inst. Netherlands*, **19**, 449.
- Van den Heuvel, E.P.J. 1976, *Structure and Evolution of Close Binary Systems*, IAU Symposium No. 73, 135.
- White, N.E. 1989, *Astr. and Ap. Rev.*, **1**, 85.
- Zahn, J.P. 1966a, *Ann. d'Ap.*, **29**, 313.
- Zahn, J.P. 1966b, *Ann. d'Ap.*, **29**, 489.
- Zahn, J.P. 1966c, *Ann. d'Ap.*, **29**, 565.
- Zahn, J.P. 1975, *Astr. and Ap.*, **41**, 329.

