

FORMATION OF VOIDS FROM NEGATIVE DENSITY PERTURBATIONS

J.C.N. de Araujo and R. Opher

Instituto Astronômico e Geofísico, USP, Brazil

RESUMEN. Se estudia la formación de huecos a partir de un espectro negativo de perturbaciones, tomando en cuenta la expansión del Universo, arrastre por fotones, enfriamiento por fotones, fotoionización, ionización colisional, enfriamiento Lyman α y la formación y enfriamiento de moléculas H_2 . Nuestros resultados predicen la existencia de regiones $\lesssim 1/10$ de la densidad promedio para regiones de masa $10^9 - 10^{10} M_\odot$.

ABSTRACT. In the present paper we study the formation of voids from a negative spectrum of perturbations taking into account the expansion of the Universe, photon-drag, photon-cooling, photoionization, collisional ionization, Lyman α cooling and the formation and cooling of H_2 molecules. Our results predict the existence of regions $\lesssim 1/10$ the average density for regions of mass $10^9 - 10^{10} M_\odot$.

Key words: CLUSTERS-GALAXIES — COSMOLOGY

I. INTRODUCTION

The formation of voids from negative density perturbations, present in the recombination era, is of appreciable interest and several investigations on the subject have been made. The previous investigations, in general, include only the effect of gravity, neglecting: photon-drag, photon cooling, photoionization, Lyman α cooling, collisional ionization and the formation and cooling of H_2 molecules. In the present investigation, we study the effect of these nongravitational physical processes in the formation of underdense regions (see also de Araujo and Opher 1990).

We use a perturbation distribution used for positive perturbations by de Araujo and Opher (1988) and postulate that a similar distribution exists for negative perturbations (de Araujo and Opher 1990):

$$\delta_i = \frac{\delta \rho}{\bar{\rho}} = - (M/M_\odot)^{-\alpha} (1+z_{\text{rec}})^{-1} \quad (1)$$

where M_\odot is the mass scale and z_{rec} is the redshift at the recombination era.

In section II we present the model and the calculations and in section III we present the conclusions and discussion.

II. MODEL AND CALCULATIONS

We assume spherical negative uniform density perturbations given by equation (1). We obtain the radius and the density of the clouds as a function of time, in particular at the present era.

We begin the calculations at $T_r = 4000\text{K}$ (beginning of recombination), (T_r - radiation temperature). We study an open universe with $\Omega_0 = 0.1$ ($\Omega_0 = \rho_0/\rho_{\text{oc}}$, ρ - present density, and ρ_{oc} - present critical density) with a Hubble constant $h = 1.0$ (in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

We take $\alpha = 1/3$ (as suggested by Gott and Rees 1975) in equation (1) with $M_\odot = 10^9$, 10^{15} and $10^{16} M_\odot$ as well as $\alpha = 1/2$ (which corresponds to a white noise Poisson perturbation spectrum) with $M_\odot = 10^{15} M_\odot$.

We also performed calculations neglecting nongravitational processes, in order to evaluate the importance of the nongravitational physical processes.

In Table 1 we present the results of the calculations for $M = 10^6 - 10^{12} M_\odot$. We note

1990RMxAA...21...38D

In general that lower the mass, M , lower the present density (i.e., lower the ratio of the present density of the perturbed region to the ambient density, ρ_p/ρ_o). We also note that the nongravitational physical processes are particularly important for small M . We also note in table 1 the present radius \bar{r}_p , of the underdense regions.

TABLE 1. Present Ratio of Density to Ambient Density, (ρ_p/ρ_o) , and Present Radius (\bar{r}_p), for Perturbed Regions of Mass M , Including Nongravitational Physical Processes (NPP) and Neglecting NPP.

M/M_\odot	With NPP		Without NPP	
	ρ_p/ρ_o	\bar{r}_p (Mpc)	ρ_p/ρ_o	\bar{r}_p (Mpc)
A. $M_o = 10^{14} M_\odot, \alpha = \frac{1}{3}$				
10^6	0.0063	0.11	0.0029	0.14
10^7	0.017	0.17	0.011	0.20
10^8	0.050	0.26	0.035	0.29
10^9	0.13	0.41	0.096	0.45
10^{10}	0.28	0.67	0.22	0.73
10^{11}	0.49	1.2	0.42	1.3
10^{12}	0.69	2.3	0.63	2.4
B. $M_o = 10^{15} M_\odot, \alpha = \frac{1}{3}$				
10^6	0.0012	0.19	0.00044	0.27
10^7	0.0048	0.26	0.0029	0.31
10^8	0.016	0.38	0.011	0.43
10^9	0.049	0.56	0.035	0.63
10^{10}	0.13	0.87	0.096	0.97
10^{11}	0.28	1.5	0.22	1.6
10^{12}	0.49	2.6	0.42	2.7
C. $M_o = 10^{16} M_\odot, \alpha = \frac{1}{3}$				
10^7	0.00082	0.47	0.00044	0.58
10^8	0.0045	0.58	0.0029	0.67
10^9	0.016	0.81	0.011	0.92
10^{10}	0.049	1.2	0.035	1.4
10^{11}	0.13	1.9	0.096	2.1
10^{12}	0.28	3.1	0.22	3.4
D. $M_o = 10^{15} M_\odot, \alpha = \frac{1}{2}$				
10^9	0.00073	2.3	0.00044	2.7
10^{10}	0.0086	2.2	0.0058	2.5
10^{11}	0.0049	2.6	0.035	2.9
10^{12}	0.20	3.5	0.15	3.9

In general, from Table 1, we produce underdense regions of present radii; \bar{r}_p , on the order of several megaparsecs. This is appreciably less than the recently discovered voids of tens of megaparsecs, which may have been produced by hierarchical clustering of smaller voids in conjunction with hierarchical clustering of galaxies (produced by the positive density perturbation spectrum).

III. CONCLUSIONS AND DISCUSSION

Our main conclusions are:

- 1) The inclusion of nongravitational physical processes (NPP) act as a "drag", impeding the formation of voids, and increasing the present density to ambient density ratio, ρ_p/ρ_o .
- 2) The ratio ρ_p/ρ_o decreases with increasing M_o .
- 3) Using Eq. 1 for $M_o/M_\odot = 10^{14}$, 10^{15} , and 10^{16} with $\alpha = 1/3$, and $M_o/M_\odot = 10^{15}$ with $\alpha = 1/2$, we produce underdense regions with present radii r_p on the order of several megaparsecs.

The distribution function of dwarf galaxies can show, in principle, the underdense regions predicted above. Crane and Saslaw (1986) compared the distribution function $f(N)$ of galaxies predicted for a quasi-relaxed gravitating system, with the observed distributions of galaxies in the Zwicky catalog. The Zwicky catalog, based on the Palomar Sky Survey plates, is reasonably complete to $m_{Zw} = 14.5$ and has a magnitude limit of $m_{Zw} = 15.5$. The Zwicky catalog is primarily sensitive to galaxies of mass $\sim 10^{11} M_\odot$. With the inauguration of the Hubble Space Telescope, and new large aperture telescopes, we can expect the obtaining of galaxy distributions with a magnitude limit ~ 21 mag and distributions of galaxies of mass $\sim 10^9 M_\odot$ can, in principle, be obtained. It would then be of interest to obtain for a given range of redshifts the distribution of dwarf galaxies of masses $\sim 10^9 M_\odot$ for small neighbor distances ($< 1^\circ$) (corresponding to fig. 2 of Crane and Saslaw (1986)).

Our results predict the existence of regions $\sim 1/10$ the average density over small neighbor distances ($< 1^\circ$), and the distribution of dwarf galaxies of masses $M \sim 10^9 M_\odot$ are sensitive to these underdense regions of mass $\sim 10^9 - 10^{10} M_\odot$.

ACKNOWLEDGEMENTS

The authors would like to thank the Brazilian agencies FAPESP and CNPq for support.

REFERENCES

- Crane, P. and Saslaw, W.C. 1986, Ap. J., 301, 1.
 de Araujo, J.C.N., and Opher, R. 1988, M.N.R.A.S., 231, 923.
 de Araujo, J.C.N., and Opher, R. 1990, Ap. J. (February).
 Gott, R.J., and Rees, M.J. 1975, Astr. Ap., 45, 365.