

PHOTOMETRY OF NOVA MUSCAE 1983

Marcos P. Diaz¹, and João E. Steiner^{2*}

1. Instituto Astronômico e Geofísico, IAG/USP, Brazil.
2. Depto. de Astrofísica, INPE/SECT, Brazil.

RESUMEN. Presentamos observaciones fotométricas de GQ Mus en su estado quieto que revelan una fuerte modulación orbital. Se discuten observaciones espectroscópicas recientes las cuales indican un período corto orbital (85 min) en fase con el período fotométrico. Basándonos en los resultados, argumentamos que el sistema pertenece a la clase de variables cataclísmicas magnetizadas.

ABSTRACT. We present photometric observations of GQ Mus at quiescence which reveal a strong orbital modulation. We also discuss recent spectroscopic observations which reveal a very short orbital period (85 min) phased with the photometric period. On the basis of the results we argue that the system belongs to the class of magnetized cataclysmic variables.

Key words: PHOTOMETRY — STARS-BINARY — STARS NOVAE

INTRODUCTION

GQ Mus is a classical nova that erupted in January, 1983 (Liller, 1983). Its outburst and decay were well covered by multiple wavelength observations from satellites (IUE, OSAT and IRAS) and in the visible, near infrared and ultraviolet from the ground. Some important properties of the system were established in the last five years: The observation of near polarization (~3%) increasing to the infrared (Cropper, 1985) and the presence of soft X-ray emission (Ögelman *et al.* 1985). At the time two possible mechanisms were proposed to explain this detection; the first was the occurrence of a hydrostatic burning process in the white dwarf surface and the second was the emission from a shock wave formed by the expanding envelope into the interstellar medium. The reddening ($E_{B-V}=0.45$) and distance (800 pc) were found by Krautter *et al.* (1984) based on the interstellar feature at 2200Å and the absolute magnitude versus decay time relation.

An increase in the degree of ionization of the ejected envelope was found and the ionizing mechanism was identified as photoionization by a central high temperature radiation source. (Krautter and Williams, 1988). An overabundance of helium, nitrogen and oxygen and probably of iron was found by Pacheco and Codina and by Krautter and Williams.

2. OBSERVATIONS AND RESULTS

The photometric observations were made in 1988 and 1989 using the 1.6m telescope at the Laboratório Nacional de Astrofísica at Brasópolis MG, Brazil. A single-channel photometer with a S11 response phototube was used to obtain the light curves with resolution of 25 seconds. These data show a remarkable high amplitude asymmetric modulation (fig. 1). No other highly significant peaks were found in a power spectrum calculated using a 6 hours light curve in the range 50-1800 seconds.

* On leave from Instituto Astronômico e Geofísico da Universidade de São Paulo.

Summary of photometric observations:

Photometric period: 0.05923 ± 0.00017 days.

T_{\min} = HJD 2,447,241.5635 \pm 0.0010

T_{\max} = 0.33 ± 0.03

Amplitude: 0.6^m (B).

Intrinsic colors: (at photometric maximum)

(U-B): -1.1

(B-V): -0.3

(V): 17.5




Fig. 1. Fast photometry of GQ Muscae.

The spectroscopic observations were made using the 2D-Frutti detector at the 4m telescope of CTIO on May, 1989. The spectra have a resolution of 3\AA , a spectral coverage from 3450\AA to 5300\AA and have a time resolution of five minutes. The spectrum of GQ Mus on May, 1989 shows a large number of lines including lines of highly ionized atoms (fig. 2). Some lines such as HeII $\lambda 4686$, OVI $\lambda 3411$ and NIII $\lambda 5270$ have a radial velocity modulation with a period which is, within the errors, equal to the photometric period (fig. 3). In these lines a periodic variation in the line profile with orbital phase is seen, a blueshifted structure appears near to the phase of maximum redshift (see the arcs in fig. 4). Some strong nebular and coronal lines formed in the nova ejecta do not show any periodic shift phased with the orbital period.




Fig. 2. Optical spectrum of GQ Muscae on May, 1989.




Fig. 3. Radial velocity curve for the OVI $\lambda 3811$ emission line.

Summary of spectroscopic observations:

Amplitude of continuum shape modulation*: $0.30^m \pm 0.15^m$

EW) HeII $\lambda 4686$: 25\AA .

EW) OVI $\lambda 3811$ + HeII $\lambda 3813$: 8\AA .

(HeII $\lambda 4686$): 400 km/s (peak).

(HeII $\lambda 4686$): 180 km/s (centroid).

(OVI $\lambda 3811$ + HeII $\lambda 3813$): 310 km/s (centroid).

ax. reddening of the continuum: $\phi = 0.0 \pm 0.1$.

ax. V (HeII $\lambda 4686$): $\phi = 0.35$.

ax. V (OVI $\lambda 3811$ + HeII $\lambda 3813$): $\phi = 0.20$.

ax. EW (HeII $\lambda 4686$): $\phi = 0.90$.

$m(3500\text{\AA}) - m(5100\text{\AA})$

II. DISCUSSION

Using the value for the period and the relations of Patterson (1984) we roughly estimate: $\dot{M} \sim 10^{-11} M_{\odot} \cdot \text{year}^{-1}$ and $M_2 \sim 0.1 M_{\odot}$ for the system at quiescence. The possibility of strong outburst sets the lower limit to the white dwarf mass down to $0.6 M_{\odot}$ (Livio and Shara 1987).

It is interesting that the large pulse fraction in the visible spectral range ($M_V = 3.3$) limits the size and temperature of a blackbody emitting region to $R^2 T = 6 \times 10^{24} \text{ K} \cdot \text{cm}^2$. Due to this condition, the modulation cannot arise from a hot white dwarf which is radiating below the Eddington limit and it should be produced in larger regions of the binary system like the heated surface of the red dwarf or the accretion stream. This fact allows us to recognize the zero phase with the photometric minimum and calculate the observed phases seen in fig. 4.

The asymmetry, the amplitude of the photometric modulation (Diaz and Steiner 1989), the modulation in the shape of the continuum and the profile variation of emission lines (Diaz and Steiner 1990) strongly suggest that the system may belong to the class of δ Her stars. Other characteristics of the system agree with this identification: the value of the orbital period, the relatively high absolute magnitude at quiescence, the low accretion rate, the intense soft X-ray luminosity and the presence of an unusually strong HeII $\lambda 4686$ emission line.




Fig. 4. A possible scenario for the emission line and continuum sources in GQ Muscae. The phases were calculated using the spectroscopic ephemeris.

IV. CONCLUSION

We have found a short orbital photometric modulation in GQ Mus and the system is proposed as a new AM Her system. GQ Mus is the nova with the shortest orbital period known, is the second object in this class with orbital period below the period gap and is the second nova with observed characteristics of polars.

REFERENCES

- Cropper, M. 1986, *M.N.R.A.S.*, **222**, 225.
 Diaz, M. P., and Steiner, J. E. 1989, *Ap. J. Lett.*, **339**, L43.
 Diaz, M. P., and Steiner, J. E. 1990, in preparation.
 Krautter, J. et al. 1984, *Astr. Ap.*, **137**, 307.
 Krautter, J., and Williams, R. E. 1988, *Messenger*, **54**, 33.
 Liller, W. 1983, *IAU Circ.*, No. 3764.
 Livio, M., and Shara, M. M. 1987, *Ap. J.*, **319**, 819.
 Ögelman, H., Beuermann, K., and Krautter, J. 1985, in *Proc. ESA Workshop: Recent Results on Cataclimic Variables* (ESA SP-236), p.177.
 Patterson, J. 1984, *Ap. J. Suppl.*, **54**, 443.

Marcos Perez Diaz e João E. Steiner: Departamento de Astrofísica, Instituto de Pesquisas Espaciais, C.P. 515, 12201 São José dos Campos (SP), Brazil.