

ON SEARCHING FOR ASTEROID FAMILIES I. HIRAYAMA FAMILIES

S. Gordillo, C. Mostaccio, R.B. Orellana,
A. Plastino, H. Tignanelli and H. Vucetich

Observatorio Astronómico
Universidad Nacional de La Plata

RESUMEN. Una nueva técnica de agrupamiento de asteroides es estudiada, basada en ideas de la taxonomía numérica. Con un software especialmente diseñado se reproducen las familias de Hirayama.

ABSTRACT: A new clustering technique based upon numerical taxonomic analysis is studied. It is shown that it faithfully reproduces Hirayama's families.

Key words: ASTEROIDS

INTRODUCTION

Hirayama (1918) was the first to determine the existence of families, i.e., groups of asteroids whose osculating elements (semi-major axis a , eccentricity e , and inclination i) have similar values.

Later on, Hirayama and other authors added new asteroids to the existing families, with the use of proper elements. Further, they determined some new families. However, the methods adopted in the search for asteroid families differ from author to author.

In the following, we show preliminary results which were obtained using methods associated with well-known numeric taxonomy procedures.

I. METHOD

Standard taxonomic techniques have been implemented (Sneath, 1978). An ad-hoc specialized software was designed to this effect. The main idea is to work within an n -dimensional "character" space. Three characters have been utilized here, namely, semimajor axis, eccentricity and inclination.

The corresponding data (associated to each asteroid), after a standard normalization procedure (Crisci and Armengol, 1983), were correlated by means of an appropriately defined similitude coefficient (distance in 3-character space (Sokal, 1961)).

Cluster analysis (Hartigan, 1975) is then invoked in order to produce the special groupings of related objects that constitute the basis of taxonomic classification schemes.

These schemes are conveniently visualized by recourse to special diagrams called "phenograms" (Willey, 1981), which in the present work describe a sample of 166 asteroids.

II. RESULTS

In order to reproduce the families found by Hirayama, we select twenty asteroids belonging to each of his original families. The pertinent orbital elements were obtained from the Ephemerides of Minor Planets (1988) and from these values, the proper elements were calculated with the tables of Brouwer and van Woerkom (1950).




Fig. 1: Reduced phenogram of 166 asteroids.




Fig. 2: Eos Family (portion of the global phenogram, fig. 1)

We applied numeric taxonomy to 166 selected asteroids. The corresponding results are arranged in the accompanying diagrams, (phenograms). Phenogram 1 is a reduced version of the full computer output, in which microstructures have been eliminated in order to provide the reader with a clear overall picture. Phenogram 2 is the section corresponding to the "Eos" family, taken as produced by our computer software. Standard numbers identifying the asteroids are given and the way in which the different groups are formed is clearly exhibited.

On the basis of the taxonomic selection criteria here employed, the original families of Hirayama are confirmed. The appropriate families are shown in the reduced phenogram ordered according to increasing values of a . Table 1, list the members of these families and give ranges of the elements. A division of the asteroids belt into three rings: inner ring (Flora), central ring (Maria), outer ring (Themis, Eos, Coronis) becomes apparent. The Phocaea separation is due to its large inclination.

TABLE 1. Hirayama Families.

Family	Members	Range
Themis	24- 62- 90-104-171-184-222-223-	a 3.08- 3.17
	268-316-379-363-431-461-468-492-	e 0.12- 0.19
	515-526-555-621	i 0.80- 2.37
Eos	221-320-339-450-513-520-529-562-	a 3.00- 3.03
	573-579-590-608-633-639-651-653-	e 0.06- 0.08
	661-669-742-766	i 9.74-10.55
Coronis	158-167-208-243-263-277-311-321-	a 2.83- 2.91
	452-462-534-658-720-761-811-832	e 0.04- 0.06
	962-975-993-1029	i 2.04- 2.27
María	170-472-575-616-652-660-695-714-	a 2.53- 2.57
	727-751-787-875-879-897-994-1158-	e 0.08- 0.12
	1160-1215-1379-1677	i 14.26-15.42
Phocaea	25-105-265-273-290-323-326-391-	a 2.30- 2.46
	502-587-654-852-914-950-1090-1108	e 0.15- 0.29
	1164-1170-1192-1316	i 18.83-26.31
Flora	8- 43-244-254-270-281-291-296-	a 2.17- 2.31
	315-341-352-364-367-376-422-428-	e 0.10- 0.18
	440-453-525-540-553-641-685-700-	i 1.11- 7.34
	703-711-736-763-770-782-800-802-	
	809-810-819-823-825-831-836-841-	
	851-913-939-956-960-963-1016-1037	
	1055-1060-1078-1088-1089-1123-	
	1130-1150-1153-1216-1225-1270-	
	1365-1377-1387-1399-1451-1619-	

IV. CONCLUSIONS

Our results suggest that automatic taxonomic techniques (where the computer does not know whether it is working with asteroids or with, say, apples) may provide the astronomer with a powerful tool for classification purposes. Indeed, we have been able in this paper to reproduce the original families of Hirayama with such an objective technique.

One is thus encouraged to process additional asteroids with these techniques, on the one hand, and to work within a larger space (that is, with other characters), on the other, in the hope of establishing new classification criteria or validating pre-existing tentative ones.

REFERENCES

- Brouwer, D. and van Woerkom, A.J.J.: 1950, *Astron. Papers, American Ephemeris*, 13, Part II.
 Crisci, J., Lopez Armengol, J.: OAS Monography N26, Washington, D.C., 1983, pgs. 51-53.
Ephemerides of Minor Planets for 1988 (Leningrad, 1988).
 Hartigan, J.A.: 1975, *Clustering Algorithms*, Wiley, Nueva York, N.Y.
 Hirayama, K.: 1918, *Proc. Phys. Math. Soc. Japan* II, 9, 354.
 Sneath, P.H.A.: 1978, *Classification of Microorganisms, Essays in Microbiology*, Morris, J.R. and Richmond, M.A. (eds), Wiley, N.Y., Cap. 9, 1.
 Sokal, R.R.: 1961, *Syst. Zool.*, 10, 70.
 Wiley, E.O.: 1981, *Phylogenetics*, Wiley, Nueva York, N.Y., XV.

ACKNOWLEDGMENT

Silvia Gordillo, Catalina Mostaccio and Horacio Tignanelli are members of CIC, Provincia de Buenos Aires, Argentina.
 Angel Plastino and Héctor Vucetich are members of CONICET, Argentina.
 Rosa Orellana and Horacio Tignanelli acknowledge partial support from IAU.

- Gordillo, C., Mostaccio, R.B., Orellana, A., Plastino, H., Tignanelli, and H. Vucetich: Observatorio Astronómico, Pádel del Bosque s/n, La Plata (1900), Argentina.