

## HIGH METALLICITY GIANT HII REGIONS

Elena Terlevich

Royal Greenwich Observatory, U. K.

### RESUMEN

Se discuten las dificultades para derivar abundancias en regiones de HII de alta metalicidad (baja excitación) debido a que en ellas, la línea de [OIII] en  $\lambda 4363 \text{ \AA}$  que es sensible a la temperatura, o es muy débil o no está presente. Usando líneas de emisión intensas sobre un amplio rango espectral (de  $\lambda 3700$  a  $\lambda 9600 \text{ \AA}$ ) se pueden determinar abundancias en dichas regiones gracias a métodos empíricos combinados con modelos de fotoionización.

### ABSTRACT

Abundances of high metallicity (low excitation) HII regions are difficult to derive, as the temperature sensitive  $\lambda 4363 \text{ \AA}$  line is very weak or absent in these objects. Empirical methods using strong emission lines covering a wide spectral range ( $\lambda 3700$  to  $\lambda 9600 \text{ \AA}$ ) combined with photoionization models provide reliable abundances for these objects.

**Key words:** GALAXIES: ABUNDANCES — GALAXIES: ISM — H II REGIONS

### 1. INTRODUCTION

Giant HII regions, which are easily seen on the discs of nearly face-on spiral galaxies, are commonly used for deriving information about the chemical composition of the interstellar medium in external galaxies. Chemical evolution models are constrained by abundance variations of different elements, and the detailed study of element ratios provides tests of nucleosynthesis and stellar evolution theories.

Giant extragalactic HII regions (GEHR) differ from Galactic ones in that they are outstandingly large and ionized by a large cluster of stars. In order to determine if standard photoionization models are adequate, we studied in detail the spatially resolved GEHR NGC 604 in M 33 (Díaz *et al.* 1987) and found that, although the excitation and measured electron temperatures vary across the nebula, neither the abundances of O, N, Ne, S, nor the empirical abundance indicators that use the intensity ratios of strong emission lines, do. These abundance indicators can therefore be used for unresolved more distant GEHR; similar results have been obtained for 30 Dor in LMC (Mathis, Chu & Peterson 1985) and NGC 5471 in M 101 (Skillman 1985).

It is known that HII regions of low excitation (as measured by the ratio  $[\text{OIII}]\lambda\lambda 4959,5007/\text{H}\beta$ ) have relatively high metal abundance. In these regions, the temperature sensitive lines required for reliable abundance determination ( $[\text{OIII}] \lambda 4363 \text{ \AA}$ ,  $[\text{NII}] \lambda 5755 \text{ \AA}$ ) are either very weak or absent. Sample spectra of a high excitation - low metallicity and a low excitation - high metallicity GEHR can be seen in Figure 1 of Terlevich, Terlevich & Franco, this volume.

Abundances of metal rich HII regions are therefore much more difficult to study. Empirical calibration of strong emission line intensity ratios with electron temperatures and abundances can be used ( $R_{23} = ([\text{OII}] + [\text{OIII}])/\text{H}\beta$  from Pagel *et al.* 1979; Alloin *et al.* 1979); the high metallicity end - where  $T_e$  is very low, elements other than O contribute to the cooling, and IR lines become important - might need to be calibrated using theoretical models (Edmunds & Pagel 1984; McCall *et al.* 1985; Dopita & Evans 1986; McGaugh 1991).

Vílchez & Pagel (1988) have developed an improved method that requires an extensive spectral coverage ( $\lambda 3727\text{--}9532 \text{ \AA}$ ) and uses simultaneous fitting of the empirical abundance parameter  $R_{23}$  and a parameter characterising the hardness of the ionizing spectrum,  $\eta = (O + /O++)/(S + /S++)$ , to an appropriate grid of

theoretical models (e.g., Stasińska 1982).  $\eta$  has the attractive properties of being almost insensitive to the ionizing parameter ( $u$ ), inversely proportional to the effective temperature  $T_{eff}$ , almost independent of reddening, almost independent of electron temperature  $T_e$ , and linear with  $\eta' = ([OII]/[OIII])/([SII]/[SIII])$ , the observed line ratios. This method has been successfully applied to determine the physical parameters of HII regions in M 33 (Díaz *et al.* 1987; Vílchez *et al.* 1988).

The results of extending this study to low excitation regions in M 51 (Díaz *et al.* 1991) using long slit spectroscopy and photoionization models are summarised in what follows.

## 2. OBSERVATIONS AND RESULTS

Long slit spectrophotometry of six GEHR in M 51 was performed at the 2.5m INT in La Palma using an Intermediate Dispersion Spectrograph with an IPCS detector covering the range  $\lambda 3600$ - $7500 \text{ \AA}$  and a CCD detector to cover  $\lambda 6200$ - $7500 \text{ \AA}$  and  $8500$ - $9800 \text{ \AA}$  giving spectral resolutions of  $4.7 \text{ \AA}$  and  $5.2 \text{ \AA}$  and spatial resolutions  $0.82$  and  $0.7 \text{ arcsec}$  respectively. [OII], [OIII] and [NII] situate all observed regions of M 51 at the low  $T_e$  end of the empirical calibration.

A second indicator related to the ionizing spectrum ( $\eta'$ ) was used in conjunction with  $R_{23}$  to select an appropriate photoionization model from a grid. GEHR emission line spectra are basically controlled by the shape of the ionizing radiation,  $u$  – the density of ionizing photons divided by particle density – and the chemical composition of the gas. In order to determine the effect of these parameters on  $R_{23}$  and  $\eta'$ , photoionization models were computed with the 1990 version of Ferland's code CLOUDY. This code, appropriate for studying HII regions of high metallicity, assumes spherical symmetry and uniform density and chemical composition for the nebula. It does not include dust. The code has the advantage though, of allowing the use of relevant values of  $u$ ; it also includes reasonable estimates of dielectronic recombination coefficients for sulphur, allowing  $S^{++}/S^{+}$  to be calculated to accuracies comparable to observational ones.

The empirical calibrator  $R_{23}$  depends mainly on the chemical composition of the gas and the  $T_{eff}$  of the ionizing stars, being almost independent of  $u$ .  $\eta'$  reflects the shape of the ionizing continuum; it depends strongly on  $T_{eff}$ , slightly on  $u$  but is independent of  $Z$ . So in the plane  $R_{23}$ - $\eta'$ , once  $u$  has been determined, one can derive  $T_{eff}$  and global metallicity values for the observed regions.  $u$  can be determined from  $S^{+}/S^{++}$  (Mathis 1985) but equally well from the observed line ratios  $[SII]\lambda\lambda 6717, 6731/[SIII]\lambda\lambda 9069, 9532$  (Díaz *et al.* 1991). Individual photoionization model fitting for each region provides line ratios, abundances, and temperatures. After this iterative process, we end up knowing  $T_{eff}$ ,  $\log u$  and  $Z$  and are ready to infer the physical properties of the nebula and its ionizing cluster. Recent evolutionary models for high metallicity giant extragalactic HII regions (García Vargas & Díaz 1993) can now be used in conjunction with the  $R_{23}$ - $\eta'$  method.

The agreement between our determination of  $H_{\alpha}$  fluxes and those from surface photometry studies (van der Hulst *et al.* 1988) indicates that our spectra are representative of the whole regions and the spectrophotometry can therefore be used to derive the number of photons required to ionize them ( $NLy$ ). The observed  $H_{\beta}$  fluxes combined with the low  $u$  found and the sizes of the regions, provide particle densities, filling factors and masses of ionized hydrogen.  $NLy$  values point to large clusters ionizing the HII regions, and therefore  $T_{eff}$  is taken to be the effective temperature of the stars that dominate the ionization. Assuming a particular IMF, one can also estimate lower limits for the mass of the ionizing clusters.

In summary, for the studied regions in M 51, we were able to determine the ionization parameter, the effective temperature of the cluster stars dominating the ionization, and the metallicity. All metallicities found are higher than solar and both ionization parameter and effective temperature are low, extending previously found trends between metallicity and ionization parameter and radiation hardness. The HII regions in M 51 are large ( $R \sim 250pc$ ), diffuse ( $n_H < 100cm^{-3}$ ) and have small filling factors. The estimated mass of ionized hydrogen is around  $10^6 M_{\odot}$  and the minimum mass of the ionizing star clusters is of the order of  $10^4 M_{\odot}$ . The small equivalent widths of  $H_{\beta}$  observed suggest contribution from older clusters.

S/O ratios are lower than solar in all the studied regions. If this effect is real and not a model artifact, it implies that the S/O ratio is lower in regions of high metallicity, which should be explained by stellar nucleosynthesis theory and chemical evolution models. N/O ratios are also different from solar in all but one of the regions, increasing with metal content, as expected if part of the nitrogen is of secondary origin.

The combination of spectrophotometric observations over a wide spectral range including the nebular [SIII] lines with adequate photoionization models, proves to be a very powerful tool for determining reliable abundances and physical conditions in low excitation regions.

These results could be improved even further by including nebular density and temperature inhomogeneities as well as dust in the photoionization models.

*Acknowledgements:* I would like to acknowledge the British and Spanish committees for allocation of telescope time (PATT and CAT) and my colleagues Angeles Díaz, Bernard Pagel, Pepe Vilchez and Mike Edmunds with whom this work has been done. It is also a pleasure to thank the organizers of this meeting and NATO for financial support.

## REFERENCES

- Alloin, D., Collin-Souffrin, S., Joly, M., & Vigroux, R. 1979, A&A, 78, 200  
 Díaz, A.I., Terlevich, E., Pagel, B., Vilchez, J.M., & Edmunds, M.G. 1987, MNRAS, 226, 19  
 Díaz, A.I., Terlevich, E., Pagel, B., Vilchez, J.M., & Edmunds, M.G. 1990, RMAA, 21, 223  
 Díaz, A.I., Terlevich, E., Vilchez, J. M., Pagel, B., & Edmunds, M.G. 1991, MNRAS, 253, 245  
 Dopita, M.A., & Evans, I.N. 1986, ApJ, 307, 431  
 Edmunds, M.G., & Pagel, B. 1984, MNRAS, 211, 107  
 García Vargas, M.L. & Díaz, A.I. 1993, ApJ, submitted  
 Mathis, J.S. 1985, ApJ, 291, 247  
 McCall, M.L., Rybski, P.M., & Shields, G.A. 1985, ApJS, 57, 1  
 McGaugh, S.S. 1991, ApJ, 380, 140  
 Pagel, B., Edmunds, M.G., Blackwell, D.E., Chun, M.S., & Smith, G. 1979, MNRAS, 189, 95  
 Stasińska, G. 1982, A&A, 48, 299  
 van der Hulst, J.M., Kennicutt, R.C., Crane, P.C., & Rots, A.H. 1988, A&A, 195, 38  
 Vilchez, J.M. & Pagel, B. 1988, MNRAS, 231, 257  
 Vilchez, J.M., Pagel, B., Díaz, A.I., Terlevich, E., & Edmunds, M.G. 1988, MNRAS, 235, 633

Elena Terlevich: Royal Greenwich Observatory, Madingley Road, Cambridge CB3 0EZ, U.K.

