

TESTING THE BIPOLAR OUTFLOW MODELS: A WELL-SHAPED BIPOLAR OUTFLOW SHELL IN MON R2

Taoling Xie¹, Paul F. Goldsmith², and Nimesh Patel

Five College Radio Astronomy Observatory, Dept. of Physics and Astronomy,
 University of Massachusetts

RESUMEN

En un mapa de CO $J = 1 - 0$, hemos identificado un cascarón bipolar delgado que delinea el lóbulo azul del flujo bipolar masivo en la región central de Mon R2. La longitud y el ancho proyectados del cascarón son respectivamente 5.3 y 2.7 pc. El espesor proyectado promedio es ~ 0.3 pc. Este cascarón angosto está notablemente bien definido y junto con los perfiles de líneas peculiares, favorece fuertemente los modelos de flujo con lóbulos casi vacíos. El modelo de flujo de Shu et al. (1991) reproduce muy bien la forma del cascarón, esto sugiere que el modelo ha logrado representar algunos de los elementos esenciales de la situación real, aunque nuestras consideraciones de la distribución de masa confirma la conclusión de un estudio anterior (Masson & Chernin 1992), que el modelo predice demasiada masa para ángulos polares pequeños ($< 14^\circ$).

ABSTRACT

In our CO $J = 1 - 0$ maps we identify an “eggplant-shaped”, thin bipolar outflow shell which outlines the extended blue lobe of the massive bipolar outflow in the central region of Mon R2. The projected length and width of the shell are about 5.3 pc and 2.7 pc respectively, and the average projected thickness of the shell is ~ 0.3 pc. This remarkably well-defined narrow shell along with the peculiar line profiles strongly favor outflow models with largely unfilled lobes. A good fit of the shape of the shell by the recent simple outflow model of Shu et al. (1991) seems to suggest that the model may have grasped some essential elements of the real situation, although our consideration of the mass distribution confirms a conclusion of a previous study (Masson & Chernin 1992) that the model predicts too much mass at small polar angles ($< 14^\circ$).

Key words: ISM: JETS AND OUTFLOWS — STARS: FORMATION
 —STARS: MASS-LOSS— STARS: PRE-MAIN SEQUENCE

1. INTRODUCTION

The bipolar outflow phenomenon has been one of the most exciting, and most intensely-studied aspects of star formation (cf. Snell 1987; Shu, Adams, & Lizano 1987). However, we may still be years away from a relatively complete understanding of the phenomenon. In fact, we cannot yet discriminate among even some very simple phenomenological outflow models. A few such models include the filled biconical lobe model by Habrit & Bertout (1986, 1990); the shell model by Moriarty-Schieven & Snell (1988); the thin paraboloidal shell model by Meyers-Rice & Lada (1991) and, more recently, the model of Shu et al. (1991, hereafter SRL) and

¹Jet Propulsion Lab, California Institute of Technology, MS 169-506, Pasadena, CA 91109 U.S.A.

²National Astronomy and Ionosphere Center, Department of Astronomy, Cornell University, Ithaca, NY 14853-6801 U.S.A.

the jet-driven outflow models of Raga & Cabrit (1993) and Masson & Chernin (1993). However, it is clear that a central element of these models is the possible existence of a molecular shell swept-up by stellar winds.

The Mon R2 bipolar outflow (Bally & Lada 1983) at a distance of 830 pc (Herbst & Racine 1976) is one of the largest in size and in mass among the two hundred or so outflows identified to date (Bally & Lane 1991). Wolf, Lada, & Bally (1990) and Meyers-Rice & Lada (1991) have presented detailed observational studies of the Mon R2 outflow and suggested possible shell structure based on the shape of line profiles, yet they do not identify limb-brightened shells.

2. OBSERVATIONS

Observations of the CO $J = 1 - 0$ transition in Mon R2 were made using the QUARRY 15-element focal plane array at the 14-m FCRAO telescope in New Salem, Massachusetts, in 1991 April through December. The map is centered on the position of an infrared star cluster, $\alpha(1950) = 06^h05^m22^s$, $\delta(1950) = -06^\circ22'25''$ (cf Beckwith et al. 1976). The spacing of the data points is 25 arcsec while the FWHM beam size of the telescope is about 45 arcsec at 115 GHz (Xie 1992; Xie, Goldsmith, & Patel 1993, hereafter XGP).

One striking feature in the maps of $^{12}\text{CO } J = 1 - 0$ emission is the appearance of an “eggplant-shaped” shell feature in several velocity channel maps with velocity close to 10.5 km s^{-1} , the centroid velocity of the core) extending from the (0,0) position to the North-West for about 20 arcmin (Xie 1992; XGP). Figure 1 presents an overlay of the high velocity gas with the shell seen in the $V_{LSR} = 10.88 \text{ km s}^{-1}$ CO channel map. The blue-shifted lobe of the bipolar outflow rests nicely within the boundary of the shell, posing the immediate possibility that we are witnessing the limb-brightened thin shell swept up by the collimated stellar wind from the central driving source of the bipolar outflow.




Fig. 1. An overlay of the shell feature (grey scale) with the high velocity gas (contours). Solid contours are for the blue-shifted emission ($V_{LSR} = 2 - 8 \text{ km s}^{-1}$) with levels 1, 2, ..., 7, 10, 13, ..., 22 $K \text{ km s}^{-1}$. The dashed contours are for the red-shifted emission ($V_{LSR} = 12 - 18 \text{ km s}^{-1}$) with levels 2, 4, ..., 26 $K \text{ km s}^{-1}$. The grey scale ranges from 1 K (white) to 14 K (black).




Fig. 2. An overlay of the modeled constant velocity locus with the velocity channel map at 10.8 km s^{-1} . The grey scale ranges from 1 K (white) to 14 K (black).

3. THE OUTFLOW SHELL

The model of SRLI and the jet-driven models of Masson & Chernin (1993) and Raga & Cabrit (1993) deal with a whole, closed outflow shell, while some other models (cf. Moriarty-Schieven & Snell 1988; Meyers-Rice & Jada 1991) have considered shells with open ends. The observed outflow shell in Mon R2 surely favors models with closed ends.

The SRLI model is particularly impressive in its simplicity and beauty in terms of mathematical construction of the physical problem, which allows a relatively easy comparison of observations with the model predictions. In a spherical polar coordinate system (r, θ, ϕ) , both the stellar wind and the ambient surrounding medium are assumed to have axial symmetry and reflection symmetry about the equatorial plane ($\theta = \pi/2$). The ram pressure force per steradian, $f(\mu) [\text{g cm s}^{-2} \text{ sr}^{-1}]$, of the wind is a function only of $\mu (= \cos \theta)$

$$f(\mu) \propto P(\mu), \quad (1)$$

and the density of the ambient material $\rho(r, \mu) [\text{g cm}^{-3} \text{ sr}^{-1}]$ can be expressed as

$$\rho(r, \mu) \propto \frac{Q(\mu)}{r^2}, \quad (2)$$

where $P(\mu)$ and $Q(\mu)$ are dimensionless functions not specified by SRLI. One approach is to let $P(\mu)$ and $Q(\mu)$ have power-law form (Masson & Chernin 1992), $P(\mu) \propto \mu^\alpha$ and $Q(\mu) \propto \mu^{-\beta}$. With these simplifications and the further assumptions that the stellar winds move out by sweeping the ambient medium into a thin shell with momentum conservation ("snowplow") and that the net mass flow along the θ direction in the thin shell is negligible, the shell velocity $v_s(\mu)$ along each radial direction ($\theta = \text{constant}$) will be

$$v_s(\mu) = v_0 \mu^\delta, \quad (3)$$

where $\delta = (\alpha + \beta)/2$ and v_0 is the shell velocity in the polar direction. Since $v_s(\mu)$ is independent of time, the shell will develop self-similarly, $r_s(\mu) = v_s(\mu)t$.

The emission from the swept-up gas in the shell is expected to show limb-brightening in some velocity

channel maps, as can be easily demonstrated by deriving the locus of constant-velocity emission from the swept-up gas.

Now how do we compare the shape of the observed shell with the prediction of the model? The simple way that XGP has taken is to parameterize 4 observable quantities of the outflow shell in 4 parameters of the model, such as δ , v_0 , inclination angle i and outflow age t , and then to determine the model parameters. The observable quantities include the maximum width, length of the shell, the maximum blue-shifted velocity along the line of sight and the projected displacement of the position where the blue-shifted velocity maximizes. We find $v_0 = 15 \text{ km s}^{-1}$, $\delta = 6$, $i = 70^\circ$ and $t = 3.5 \times 10^5 \text{ yrs}$. This inclination angle is consistent with the 66 determined by Meyers-Rice & Lada (1991). Figure 2 presents an overlay of the modeled constant velocity loci with the velocity channel map at 10.8 km s^{-1} . We see that the SRL model provides a remarkable fit to the shape of the observed outflow shell.

4. DISCUSSION

Besides the shape of the shell, another aspect of the model is the mass distribution in the shell, which depends on α and β respectively. Masson & Chernin (1992) investigated this aspect by studying the predicted line profile averaged over the whole outflow, and concluded that the SRL model predicts too much mass at small polar angles. XGP presented a different approach. They parameterized the column density sampled by the telescope beam as a function of the centroid velocity of the blue-shifted shell along the outflow axis. If the ^{12}CO emission from the shell is assumed to be optically-thin, then this function corresponds to the integrated intensity of ^{12}CO as a function of the shell velocity along the outflow axis. XGP found that while the SRL model in its present form does lead to far more mass at very small angles than observed, it seems to provide satisfactory explanations to the mass distribution as a function of velocity over a large range of polar angles. Being encouraged by this and the remarkable fit to the shape of the outflow shell, XGP argue that the SRL model may have grasped the essential physics of the phenomenon and the discrepancy at very small polar angle between the model and the data may be due to possible complications such as inhomogeneity, clumpiness, etc. which are not properly characterized by the over-simplified model at present.

One thing worth mentioning is that we have only treated the blue-shifted lobe so far. The outflow actually has considerable structure and complications (Meyers-Rice & Lada 1991; Xie 1992; XGP). Two of these complications seen in Figure 1 are the "bending" of the red-shifted lobe and the serious overlapping of red- and blue-shifted high velocity gas close to the outflow center which is not expected for the estimated inclination angle of $\sim 70^\circ$. Xie (1992) suggests that most of these can be understood in that the central core and the bipolar outflow are located on a large expanding molecular shell which now dominates the overall giant molecular cloud and triggered new generations of star formation (Xie & Goldsmith 1994). While the blue lobe of the outflow is developing into the relatively homogeneous medium which has not yet been swept-up by the large scale expansion, the red lobe is developing towards the cavity which is largely cleared out of molecular material and the red lobe is "squashed" so that it is forced to develop sideways.

TX acknowledges E.E. Mendoza and family for their kind hospitality, and sincerely thanks A. Arellano Ferro and colleagues for the well organized symposium and also for providing partial support for his stay at Mexico City.

REFERENCES

- Bally, J., & Lada, C.J. 1983, *ApJ*, 265, 824
- Bally, J., & Lane, A.P. 1991, in *The Physics of Star Formation and Early Stellar Evolution*, ed. C.J. Lada & N.D. Kylafis (Dordrecht: Kluwer), p. 471
- Beckwith, S., Evans II, N.J., Becklin, E.E., & Neugebauer, G. 1976, *ApJ*, 208, 390
- Cabrit, S., & Bertout, C. 1986, *ApJ*, 307, 313
- _____. 1990, *ApJ*, 348, 530
- Herbst, W., & Racine, R. 1976, *AJ*, 81, 840
- Masson, C.R., & Chernin, L.M. 1992, *ApJ*, 387, L47
- _____. 1993, *ApJ*, 414, 230
- Meyers-Rice, B.A., & Lada, C.J. 1991, *ApJ*, 368, 445
- Moriarty-Schieven, G.H., & Snell, R.L. 1988, *ApJ*, 333, 316
- Raga, A., & Cabrit, S. 1993, *A&A*, 278, 267
- Shu, F.H., Adams, F.C., & Lizano, S. 1987, *ARA&A*, 25, 23

1994RMxAA...29...63X

- Shu, F.H., Ruden, S.P., Lada, C.J., & Lizano, S. 1991, ApJ, 370, L31 (SRLL)
Snell, R.L. 1987, Star Forming Regions, ed. M. Peimbert & J. Jugaku (Dordrecht: Reidel), p. 213
Wolf, G.A., Lada, C.J., & Bally, J. 1990, AJ, 100, 1892
Xie, T. 1992, Ph.D. thesis, University of Massachusetts at Amherst
Xie, T., Goldsmith, P.F., & Patel, N. 1993, ApJ, 419, L33 (XGP)
Xie, T., & Goldsmith, P.F. 1994, ApJ Letters, July 10 issue

P.F. Goldsmith, N. Patel and T. Xie: Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Dr., Pasadena, CA 91109 U.S.A. e-mail: tao@koala.jpl.nasa.gov.

