

## RESEARCH NOTE: $\beta$ AND $\alpha$ RECOMBINATION LINES FROM A REGION AROUND THE GALACTIC CENTER

I.N. Azcárate<sup>1</sup>

Instituto Argentino de Radioastronomía

and

J.C. Cersosimo

University of Puerto Rico, Dept. of Physics and  
Electronics, Humacao Univ. College

*Received 1994 April 11*

### RESUMEN

Se realizaron observaciones de las líneas de recombinación  $H159\alpha$  ( $\nu = 1620.672$  MHz) y  $H200\beta$  ( $\nu = 1619.690$  MHz) en nueve posiciones alrededor de la región del centro galáctico (separadas entre sí por  $0.5^\circ$ ), utilizando la antena de 30-m del Instituto Argentino de Radioastronomía. El propósito de las observaciones fue estimar el apartamiento de las condiciones de Equilibrio Termodinámico Local (ETL) en esa región. Sólo en dos de las nueve posiciones, se pudo detectar la línea  $H200\beta$  por encima del nivel de ruido. Para esas dos posiciones, se determinó el cociente:  $P(H200\beta)/P(H159\alpha)$ , con el objeto de verificar el apartamiento de las condiciones de ETL.

### ABSTRACT

$H159\alpha$  ( $\nu = 1620.672$  MHz) and  $H200\beta$  ( $\nu = 1619.690$  MHz) recombination line observations of nine positions around the galactic center region (separated by  $0.5^\circ$  from each other) were made with the 30-m antenna of the Instituto Argentino de Radioastronomía. The observations were made with the purpose of checking the departure from the Local Thermodynamic Equilibrium (LTE) conditions in this region. Only in two positions the  $H200\beta$  line was detected above the noise-level. The ratio  $P(H200\beta)/P(H159\alpha)$  was determined for these two positions, in order to check the departure from LTE conditions.

*Key words:* **GALAXY-CENTER — H II REGIONS**

### 1. INTRODUCTION

Important physical information can be obtained from radio recombination line observations of H II regions, such as electron density, emission measure, and electron temperature of the nebulae. The importance of stimulated emission effects in the observed radio recombination lines that strengthen the intensity of the lines was first pointed out by Goldberg (1966). He was the first to suggest that the inten-

sity of radio recombination lines can be essentially dependent on the non-equilibrium of the population of highly excited levels (non-LTE effect). Since then, the possible presence of stimulated emission has been taken into account for the interpretation of recombination line observations (Dupree & Goldberg 1969; Brocklehurst & Seaton 1972; Brown, Lockman, & Knapp 1978). The stimulated emission produces departures from LTE conditions in the ionized gas.

These non-LTE effects can be investigated by observations of  $\alpha$  and  $\beta$  transitions at about the same frequency (Dupree & Goldberg 1970). In the case of  $H159\alpha$  and  $H200\beta$  the  $P(H200\beta)/P(H159\alpha)$ , integrated intensity line ratio, is  $\simeq 0.28$  for LTE conditions.

<sup>1</sup> Member of the Carrera del Investigador Científico y Tecnológico del Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina.

To investigate the presence of non-LTE effects in a region around the galactic center, we have made observations of both lines in nine positions in this zone (separated  $0.5^\circ$  from each other). The frequencies of the lines are: 1620.672 MHz for the  $\alpha$  and 1619.690 MHz for the  $\beta$  transition.

## 2. THE OBSERVATIONS

The observations were made with the 30-m antenna of the Instituto Argentino de Radioastronomía. The receiver was a room temperature GaAs FET amplifier. The system temperature on cold sky was 89 K. The aperture efficiency was  $\eta_a = 0.45$ . The conversion factor from antenna temperature to flux densities for the system is 8.7 Jy/K. The antenna beam at 18-cm was  $29'$ . The spectral analysis was performed with a filter-bank of 84 channels of 75 kHz width each; that allowed a velocity resolution of  $13.9 \text{ km s}^{-1}$ . The frequency switching mode was used to make the observations. The local oscillator was switched by an amount less than the observing bandwidth, to obtain two independent spectra, which were then averaged. Third and fourth order polynomials were used to correct baseline effects. The integration time for each position was 4–5 hr, which resulted in an r.m.s. of 7 mK. The integration time for the observation of the galactic center itself ( $\ell = 0.0^\circ$ ,  $b = 0.0^\circ$ ) was 6 1/2 hr.

## 3. RESULTS

The following points were observed:  $\ell = -0.5^\circ$ ,  $b = +0.5^\circ$ ;  $\ell = -0.5^\circ$ ,  $b = 0.0^\circ$ ;  $\ell = -0.5^\circ$ ,  $b = -0.5^\circ$ ;  $\ell = 0.0^\circ$ ,  $b = +0.5^\circ$ ;  $\ell = 0.0^\circ$ ,  $b = 0.0^\circ$ ;  $\ell = 0.0^\circ$ ,  $b = -0.5^\circ$ ;  $\ell = +0.5^\circ$ ,  $b = +0.5^\circ$ ;  $\ell = +0.5^\circ$ ,  $b = +0.0^\circ$ ;  $\ell = +0.5^\circ$ ,  $b = -0.5^\circ$ . Only in the galactic center itself (G0.0+0.0), and the point G0.5–0.5 the H200 $\beta$  line was detected above the noise level. We show in Table 1 the observed line parameters for these two cases. In this table,  $T_L$  is the line temperature,  $\Delta v$  is the linewidth,  $V_{lsr}$  is the central velocity of the line, and  $T_c$  is the continuum temperature. We discuss below the results.




Fig. 1. Profile corresponding to the G0.0–0.0 position. In the ordinate axis, the antenna temperature is given in mK. In the abscissa axis the frequency is given in MHz. The origin of this last axis corresponds to the laboratory frequency of the H159 $\alpha$  line. The LSR velocities are given in the upper part of each peak. The same indications are valid for Figure 2.

### 3.1. G0.0+0.0

The profile corresponding to this position is shown in Figure 1. The  $P(\text{H}200\beta)/P(\text{H}159\alpha)$  ratio is  $0.24 \pm 0.04$ , where  $P(\text{H}200\beta)/P(\text{H}159\alpha)$  is the integrated intensity line ratio. We could interpret this result as an indicator that the LTE departures are not significant. That could be explained by the presence of overlapping low-density H II regions along the line of sight. In these low-density H II regions and with this large beam ( $29'$ ) we would not have observable non-LTE effects (Shaver 1980a, b). To have  $\beta/\alpha$  ratios close to 0.28 is not enough to decide that the LTE departures are not significant, since the  $\beta$  and  $\alpha$  lines can be out of the LTE in the same direction and the ratio still be 0.28. The derived electron temperature from the H159 $\alpha$  observation is  $6960 \pm 400 \text{ K}$ .

TABLE 1

OBSERVED LINE PARAMETERS

Region	$\alpha$ (1950)	$\delta$	Line	$T_L$ (K)	$\Delta v$ ( $\text{km s}^{-1}$ )	$V_{lsr}$ ( $\text{km s}^{-1}$ )	$T_c$ (K)
G0.0+0.0	265.604	–28.927	{H159 $\alpha$	0.45	$50 \pm 3$	$+8 \pm 3$	$61 \pm 3$
			{H200 $\beta$	0.11	$50 \pm 5$	$-2 \pm 3$	
G0.5–0.5	266.389	–28.762	{H159 $\alpha$	0.29	$40 \pm 3$	$+8 \pm 3$	$17 \pm 2$
			{H200 $\beta$	0.09	$26 \pm 3$	$-2 \pm 3$	

### 3.2. G0.5–0.5

We show in Figure 2 the profile corresponding to galactic coordinates  $\ell = 0.5^\circ$ ,  $b = -0.5^\circ$ . The  $\beta/\alpha$  ratio is  $0.20 \pm 0.04$  which is an indication of the presence of non-LTE effects. These effects can be attributed to stimulated emission caused by the strong non-thermal galactic continuum radiation. In this case, the number of H II regions (probably most of them of low density) that appear in the line of sight is almost certainly much lower than in the direction of the galactic center. On the other hand, the low  $\beta/\alpha$  ratio could not be explained by pressure broadening effects, since the broadening in the  $\beta$  line is not larger than that of the  $\alpha$  line. Therefore, what we see is most probably  $\alpha$  and  $\beta$  emission in the region G0.5–0.5 itself, stimulated by the non-thermal galactic continuum radiation.

The derived electron temperature from the H159 $\alpha$  observation at this position is  $4100 \pm 300$  K. For the other seven observed positions, we can give an upper limit to the H200 $\beta$  line intensity of 25 mK.




Fig. 2. Profile corresponding to the G0.5–0.5 position.

### 4. CONCLUSIONS

In this basically observational work, we have carried out H159 $\alpha$  and H200 $\beta$  recombination line obser-

vations in a region around the galactic center (nine positions) with the 30-m antenna of the IAR, with a HPBW of 29'. Due probably to the relatively low sensitivity of the receiver ( $T_{sys} \simeq 89$  K), only in two positions the H200 $\beta$  line was detected. For these positions we obtained  $\beta/\alpha$  ratios which in one of the positions (G0.5–0.5) seem to indicate the presence of non-LTE effects, caused most probably by the non-thermal galactic continuum radiation. For this position, the low  $\beta/\alpha$  could not be explained by the presence of pressure broadening effects.

For the other position, corresponding precisely to the galactic center (G0.0+0.0) there does not seem to be stimulated emission. The explanation for this fact could be the presence of a superposition of low-density H II regions, in which there would not be observable non-LTE effects for observations made with the large beam used by us, of 29' (Shaver 1980a,b). We do not have observational evidence of the presence of pressure broadening effects in this case.

The upper limits for the positions with no positive  $\beta$  detection are 25 mK for the intensity of this recombination line. More sensitive observations would be necessary to detect the H200 $\beta$  line and test the possible departures from LTE conditions in the other positions.

We wish to thank the technical staff of the Instituto Argentino de Radioastronomía, for their support during the observations. We thank the referee for the very useful comments to improve the paper.

### REFERENCES

- Brocklehurst, M., & Seaton, M.J. 1972, MNRAS, 157, 179
- Brown, R.L., Lockman, F.J., & Knapp, G.R. 1978, ARA&A, 16, 445
- Dupree, A.K., & Goldberg, L. 1969, ApJ, 158 L49
- \_\_\_\_\_. 1970, ARA&A, 8, 231
- Goldberg, L. 1966, ApJ, 144, 225
- Shaver, P.A. 1980a, A&A, 90, 34
- \_\_\_\_\_. 1980b, A&A, 91, 279

Ismael N. Azcárate: Instituto Argentino de Radioastronomía, Casilla de Correo No. 5, Villa Elisa (1894), Prov. de Buenos Aires, Argentina. (azcarate@irma.edu.ar).

J.C. Cersosimo: University of Puerto Rico, Dept. of Physics and Electronics, Humacao University College, PR 00791, Puerto Rico.

