

mixed with primary contributions leading to qualitative and/or quantitative imprecise detections.

A variety of methods are being used to search for different NG signals in CMB data because one does not expect that a single statistical tool can be able to identify all possible forms of Gaussian deviations. Using two directional large-angle NG indicators based on skewness and kurtosis statistical momenta of patches of the CMB sphere, we analyze the three nearly full sky foreground-cleaned *Planck* maps: SMICA, NILC, and SEVEM.

Our results show that these foreground-cleaned *Planck* maps exhibit different levels of NG at large angles, depending on the cut-sky mask used (all of them released by the *Planck collaboration*). The separation component minimum mask, termed M82, and the U73 mask appear to be equally efficient to Gaussianize all these CMB *Planck* maps.

¹ Observatório Nacional, MCTI.

CENTRAL DOMINANT GALAXIES AND THE EVOLUTION OF THEIR HOST CLUSTERS
C. A. Caretta¹, H. Andernach¹, J. J. Trejo-Alonso¹,
J. de Anda-Suárez², H. Santoyo-Ruiz², M. A.
Muñiz-Torres², and C. Hernández-Aguayo²

We have studied a sample of 50 galaxy clusters, all with more than 100 spectroscopically confirmed member galaxies, from all Bautz-Morgan types, in order to determine the importance of the brightest cluster members (BCMs) and their relation to the structure and dynamical state of their host clusters. Strict tests for membership and for the presence of substructures were applied. Near-infrared data were used for photometry and astrometry, which allowed us to rank the member galaxies in order of their stellar mass content. The large spectroscopic sampling, the broad range of morphologies, masses and intra-cluster medium properties, beyond the improved analyses for membership and substructuring, make this sample an especially suitable reference of nearby optical clusters ($0.005 < z < 0.150$) for the study of cluster evolution and environment effects on member galaxies. Only 35% of the clusters revealed to be unimodal, 50% to be substructured and other 15% to be multi-modal. Also, for about 20% of the sample, the Central Dominant Galaxy (CDG) of the main structure is not the first-ranked BCM, but the CDG of a substructure. More massive clusters present more than one dominant galaxy, while the

less massive ones present only one, if any. This correlation goes in the sense that most of the evolution of CDGs occurs in groups that are doomed to merge and form clusters.

¹ Departamento de Astronomía, DCNE-CGT, Universidad de Guanajuato; Callejón de Jalisco, S/N, Col. Valenciana, 36240, Guanajuato, Gto., Mexico (caretta,heinz.josue@astro.ugto.mx).

² Departamento de Física, DCI-CLE, Universidad de Guanajuato; Loma del Bosque, 103, Col. Lomas del Campestre, 37150, León, Gto., Mexico.

NEW CATALOGUES OF SUPERCLUSTERS OF ABELL/ACO GALAXY CLUSTERS UP TO
 $z \sim 0.15$

M. Chow Martínez¹, H. Andernach¹, and C. A. Caretta¹

We present two new catalogues of superclusters of galaxies within a redshift of $z = 0.15$, constructed using the Abell/ACO cluster redshift compilation by one of us (H.A.). One is the all-sky Main SuperCluster Catalogue (MSCC), based on 3410 A-clusters (92% with spectroscopic redshifts), containing 601 superclusters with multiplicities from 2 to 42; the other is the Southern ($\delta < -17^\circ$) SuperCluster Catalogue (SSCC) based on 1227 A-clusters and 1177 S-clusters (90% with spectroscopic redshifts), containing 425 superclusters with multiplicities from 2 to 39. These are currently the deepest all-sky supercluster catalogues based on optical data. By comparing both catalogues, we found the following effects (expressed as percentages of the total number of superclusters in SSCC): new superclusters with S-clusters around A-cluster cores (12.6%), the formation of bridges of S-clusters between A-clusters (1.2%), and the addition of new superclusters formed by S-clusters only (25.4%). We determined morphological parameters for the superclusters, based on ellipsoid fits and Minkowski functionals, obtaining that 39% of the rich (multiplicity >5) superclusters are prolate ellipsoids and 61% are oblate ones. The cumulative multiplicity functions of both catalogues follow very closely a power law with an exponent of -2.0 for MSCC and -1.9 for SSCC. This power law is clearly inconsistent with the same function we derived for supercluster catalogues based on simulated samples of randomly distributed clusters. It is also inconsistent, though less significantly, with similar analyses we applied to the Bolshoi cosmological sim-